arXiv:1204.2337v2 [quant-ph] 30 Jul 2012

General properties of the maximized entanglement of indirectly interacting two spins

Tomotaka Kuwahara[1]

[1] *Department of Physics, The University of Tokyo, Komaba, Meguro, Tokyo 153-8505*

We investigate the thermal entanglement of indirectly interacting two spins through other spins, that is, two spins at the ends of a spin chain. We maximize it by tuning the local fields on the two spins to obtain the maximized entanglement. We prove that if the two spins are separated by two sites or more, there is a critical temperature above which the maximized entanglement vanishes. We numerically calculate the maximized entanglement in three-spin chains and four-spin chains. We discover that the maximizing local fields on the spins 1 and 2 have asymmetric forms, which implies that the asymmetry of the two spins essentially contributes to the entanglement enhancement. In the three-spin chains, we explain this enhancement due to the asymmetry qualitatively and quantitatively in terms of the magnons. In XX and XY four-spin chains, we find that the critical temperature shows qualitatively different behavior depending on the conservation of the angular momentum in the z direction.

I. INTRODUCTION

The quantum entanglement [1, 2] is an essential resource for quantum information processing [3, 4] and hence it is very important to generate the entanglement efficiently. The entanglement in thermal equilibrium (the thermal entanglement [5]) has been researched energetically [6–10] because the destruction of the entanglement by thermal fluctuation extremely restricts the entanglement generation. It is therefore of great importance to know how much entanglement can be generated in thermal equilibrium [11–13]. We thereby consider here the maximum value of the entanglement in thermal equilibrium. In Ref. [14], it was shown for two spins interacting directly with each other, how much entanglement we can enhance between the spins by tuning the local fields on the spins. In the present paper, we will extend the maximization problem to multipartite spin systems; we study the enhancement of the bipartite entanglement due to the local fields in the system where the two spins *indirectly* interact with each other via other spins. The main research targets are the following issues:

- Is it possible to generate the entanglement at high temperatures in the indirectly interacting spins?
- What are the main factors which make it possible for the local fields to enhance the entanglement?

For the directly interacting two spins [14], the answers for these questions have been obtained: the maximized entanglement decays as $1/(T\ln T)$ as the temperature increases; the entanglement enhancement by the local fields is brought about by the suppression of the thermal fluctuation, or in other words, the increase of the purity. Our purpose in the present paper is to answer these questions qualitatively and quantitatively for indirectly interacting spins.

The possibility of the enhancement of the entanglement with the external fields has been shown in many papers [6, 11–13, 15–17]. However, there have been few reports about the entanglement maximization of specific two spins. Moreover, little is known about the mechanism of the entanglement enhancement due to the external fields or the condition for the indirect interaction to generate the entanglement. By making these problems clear, we will be able to know the fundamental limits of the entanglement generation in thermal equilibrium.

In our research, we define the maximized entanglement as the maximum value of the entanglement between the particular two spins under the condition that we can arbitrarily tune the local fields only on these two spins. We regard the other parameters such as interaction parameters as fixed. If the maximized entanglement is equal to zero, the entanglement generation is impossible for any values of the local fields. The forms of the maximizing local fields reflect basic properties of the entanglement enhancement, and thereby help us solve the above two problems.

We study general properties of the maximized entanglement between the focused two spins which are connected by a spin chain. Our main results are the following two:

- At high temperatures, we prove that the maximized entanglement is always equal to zero between the two spins which are separated by two or more spins. In other words, above a critical temperature $\tilde{T}_{\rm c}$, we can never generate the entanglement between spins far apart for any values of the local fields. It is well known [5] that the entanglement vanishes above a certain temperature because of thermal disturbance, but if we can control the local fields on the focused spins, the increase of the purity owing to the local fields may suppress the effect of thermal fluctuations. Indeed, in two and three spin chains, the non-zero entanglement can be achieved at any temperatures by increasing the local fields of the two spins. A strong local field causes the decoupling of the two spins, bringing them close to the product state, but the suppression of the purity is more dominant in two



FIG. 1: Schematic picture of the spin chains. We define the spins 1 and N as the focused spins and the spins which mediate the indirect interaction between the spins 1 and N as the media spins. We assume that we can tune the local fields on the spins 1 and N, while the other fields $\{h_i^z\}_{i=2}^{N-1}$ are arbitrary but fixed.

and three spin chains. On the other hand, in more than three spins, the decoupling by the local fields is more dominant than the suppression of the purity, and hence we cannot keep non-zero entanglement for any values of the local fields above a critical temperature.

- The main factors of the entanglement enhancement due to the local fields are not only the increase of the purity but also their effect on the indirect interaction. To be more specific, the external fields affect the magnons which mediate the indirect interaction. The form of the maximizing local fields depends on this effect and has asymmetric forms in a particular parameter region.

We show these results analytically and numerically. This paper consists of the following sections: in Section II, we state the problem specifically and give some definitions; in Section III, we give a general theorem on the entanglement enhancement which can be applied to any spin chains; in Section IV, we show the numerical and analytical results on the maximization of the entanglement in three-spin chains; in Section V, we show the numerical and analytical results on the maximization of the entanglement in four-spin chains; and in Section VI, discussion concludes the paper.

II. STATEMENT OF THE PROBLEM

First, we formulate the framework of the entanglement maximization problem and describe conditions. We consider a general XYZ N-spin chain with external fields in the z direction. The most general form of the Hamiltonian of this system is given as follows:

$$H_{\mathrm{tot}} = \sum_{i=1}^{N-1} (J_i^x \sigma_i^x \sigma_{i+1}^x + J_i^y \sigma_i^y \sigma_{i+1}^y + J_i^z \sigma_i^z \sigma_{i+1}^z) + \sum_{i=1}^{N} h_i^z \sigma_i^z, \tag{1}$$

where $\{\sigma_i^i\}_{i=x,y,z}$ are the Pauli matrices and we adopt the free boundary conditions.

We hereafter consider the entanglement between the spins 1 and N at the ends of the chain. We refer to these two spins as the focused spins and to the external fields h_1^z and h_N^z on the two spins as the local fields. We refer to the other spins $2 \le i \le N-1$ as the media spins (Fig. 1). The basic problem that we are going to solve is to maximize the thermal entanglement between the focused spins 1 and N by tuning the local fields h_1^z and h_N^z at a fixed temperature. We also fix all the other parameters, namely, $\{J_i^x, J_i^y, J_i^z\}$ for $1 \le i \le N-1$ and $\{h_i^z\}$ for $2 \le i \le N-1$. We refer to the maximizing values of the local fields h_1^z and h_N^z as $h_{1\mathrm{op}}$ and $h_{N\mathrm{op}}$.

Note that the maximizing local fields $h_{1\mathrm{op}}$ and $h_{N\mathrm{op}}$ generally depend on the temperature T, or on the inverse temperature $\beta = 1/(kT)$ with k the Boltzmann constant. This is because we tune the local fields at a fixed temperature β. Let us then define the high-temperature limit, in which we mostly develop the argument. In our high-temperature limit, we make $\beta||H_{\mathrm{tot}} - h_1^z\sigma_1^z - h_N^z\sigma_N^z||$ tend to zero, where $||\cdots||$ denotes the matrix norm. On the other hand, we let the maximizing local fields depend on β as we take the limit $\beta \to 0$. Hence, $\beta h_{1\mathrm{op}}$ and $\beta h_{N\mathrm{op}}$ may even diverge in our high-temperature limit.

The density matrix of the total system in thermal equilibrium is

$$\rho_{\mathrm{tot}} = \frac{e^{-\beta H_{\mathrm{tot}}}}{Z_{\mathrm{tot}}}, \tag{2}$$

where $Z_{\rm tot} = {\rm tr}(e^{-\beta H_{\rm tot}})$ is the partition function. The density matrix of the focused spins 1 and N is

$$\rho_{1N} = {\rm tr}_{1N}\rho_{\rm tot}, \tag{3}$$

where ${\rm tr}_{1N}$ denotes the trace operation on the spins *except* the focused spins 1 and N. For the present system (1), the general form of the density matrix ρ_{1N} is given by:

$$\rho_{1N} = \begin{pmatrix} p_{\uparrow\uparrow} & 0 & 0 & F_2 \\ 0 & p_{\uparrow\downarrow} & F_1 & 0 \\ 0 & F_1 & p_{\downarrow\uparrow} & 0 \\ F_2 & 0 & 0 & p_{\downarrow\downarrow} \end{pmatrix} \tag{4}$$

in the basis of the eigenstates of $\sigma_1^z \otimes \sigma_N^z$, where $p_{\uparrow\uparrow}$, $p_{\uparrow\downarrow}$, $p_{\downarrow\uparrow}$, $p_{\downarrow\downarrow}$, F_1 and F_2 are real numbers. We have $F_2 = 0$ when $J_i^x = J_i^y$, for $1 \le i \le N-1$, in particular.

In order to quantify the entanglement, we here adopt the concurrence [18] as an entanglement measure. The concurrence $C(\rho_{1N})$ is defined as follows:

$$C(\rho_{1N}) \equiv \max(\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4, 0), \tag{5}$$

where $\{\lambda_n\}_{n=1}^4$ are the eigenvalues of the matrix

$$\sqrt{\rho_{1N}(\sigma_1^y \otimes \sigma_N^y)\rho_{1N}^*(\sigma_1^y \otimes \sigma_N^y)} \tag{6}$$

in the non-ascending order $\lambda_1 \ge \lambda_2 \ge \lambda_3 \ge \lambda_4$. For density matrices of the form (4), the concurrence $C(\rho_{1N})$ is reduced to the simpler form

$$C(\rho_{1N}) = 2\max(|F_1| - \sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}}, |F_2| - \sqrt{p_{\uparrow\downarrow}p_{\downarrow\uparrow}}, 0). \tag{7}$$

Then, the necessary and sufficient condition for the existence of the entanglement is given by

$$\max(|F_1| - \sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}}, |F_2| - \sqrt{p_{\uparrow\downarrow}p_{\downarrow\uparrow}}) > 0. \tag{8}$$

Thus, the present entanglement optimization problem for the spin pair $(1, N)$ is equivalent to finding the values of $\{h_1^z, h_N^z\}$ which maximize $C(\rho_{1N})$ with the parameters $\{J_i^x, J_i^y, J_i^z\}$ $(1 \le i \le N-1)$, $\{h_i^z\}$ $(2 \le i \le N-1)$, and β all fixed.

III. GENERAL THEOREMS ON ENTANGLEMENT GENERATION

In the previous section, we formulated the entanglement maximization problem. In fact, there are cases in which we cannot generate the entanglement for any values of the local fields at all. In such cases, it is necessary to distinguish whether the entanglement is exactly equal to zero or rapidly approaches to zero. In the present section, we introduce a general theorem on a necessary condition to generate the entanglement by optimizing the local fields. In other words, we give a sufficient condition for the entanglement to be zero exactly for any values of the local fields.

Theorem 1. Let us consider the XYZ chain (1) with $\{J_1^x, J_1^y, J_1^z\} = \{J_{N-1}^x, J_{N-1}^y, J_{N-1}^z\} = \{J^x, J^y, J^z\}$. We tune the local fields h_1^z and h_N^z, while the media fields $\{h_i^z\}_{i=2}^{N-1}$ are arbitrary but fixed. There is a critical temperature above which the maximized entanglement between the focused spins 1 and N is exactly zero if they are separated by two or more spins ($N \ge 4$). In other words, we cannot generate the entanglement for any values of the local fields above this temperature for $N \ge 4$.

Comments. This critical temperature yields a stronger restriction than the well-known ones [5], which discuss the entanglement disappearance under a fixed Hamiltonian. Let us denote this critical temperature as $\tilde{T}_{\rm c}$. We say $\tilde{T}_{\rm c} = \infty$ if we can generate the entanglement at any temperatures, whereas we say $\tilde{T}_{\rm c} = 0$ if the maximized entanglement is equal to zero even in the ground state.

Proof. In order to prove this theorem, it is enough to show

$$\max(F_1^2 - p_{\uparrow\uparrow}p_{\downarrow\downarrow}, F_2^2 - p_{\uparrow\downarrow}p_{\downarrow\uparrow}) < 0 \tag{9}$$

after the maximization of the left-hand side with respect to h_1^z and h_N^z in the high temperature limit $\beta \to 0$, where $\{F_1, F_2, p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$ are the elements of the density matrix defined in Eq. (4). Then, Eq. (7) yields the exactly zero concurrence in the limit $\beta \to 0$. Since the system (1) has a finite number of degrees of freedom, the elements

$\{F_1, F_2, p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$ must be analytic as a function of β. Therefore, there can be a finite value of β at which $\max(F_1^2 - p_{\uparrow\uparrow}p_{\downarrow\downarrow}, F_2^2 - p_{\uparrow\downarrow}p_{\downarrow\uparrow}) = 0$ after the maximization. This gives the critical temperature. Note that the elements of ρ_{1N} here are functions of $h_{1\mathrm{op}}(\beta)$, $h_{N\mathrm{op}}(\beta)$ and β.

Now, let us define the exponents κ_1 and κ_N as

$$h_{1\mathrm{op}} = O(\beta^{-\kappa_1}) \text{ and } h_{N\mathrm{op}} = O(\beta^{-\kappa_N}) \tag{10}$$

in the limit $\beta \to 0$, where κ_1 and κ_N are real numbers. We estimate the order of each element of $\{F_1^2, F_2^2\}$ and $\{p_{\uparrow\uparrow}p_{\downarrow\downarrow}, p_{\uparrow\downarrow}p_{\downarrow\uparrow}\}$ in the following three cases:

- Case (a): $\kappa_1 < 1$ and $\kappa_N < 1$.
- Case (b): $\kappa_1 \geq \kappa_N$, $\kappa_1 \geq 1$ and $\kappa_N > 0$; or $\kappa_N \geq \kappa_1$, $\kappa_1 > 0$ and $\kappa_N \geq 1$.
- Case (c): $\kappa_1 \geq 1$ and $\kappa_N \leq 0$; or $\kappa_1 \leq 0$ and $\kappa_N \geq 1$.

Notice that the three cases cover the entire space of κ_1 and κ_N.

Case (a). In this case, we can prove the inequality (9) by utilizing a necessary condition for the existence of the entanglement [19], which is

$$\mathrm{tr}\,\rho_{1N}^2 \geq \frac{1}{3}. \tag{11}$$

In the case (a), $\beta h_{1\mathrm{op}}$ and $\beta h_{N\mathrm{op}}$ are of order $\beta^{1-\kappa_1}$ and $\beta^{1-\kappa_N}$, respectively, and approach to zero in the high temperature limit $\beta \to 0$. The density matrix ρ_{1N} becomes proportional to the identity matrix because then $\beta||H_{\mathrm{tot}}|| \to 0$; hence we have

$$\lim_{\beta \to 0} \mathrm{tr}\,\rho_{1N}^2 = \frac{1}{4}. \tag{12}$$

Therefore, in the case(a), the entanglement between the spins 1 and N is exactly zero in the high-temperature limit.

Case (b). To simplify the problem, we consider the case of $h_{1\mathrm{op}}, h_{N\mathrm{op}} > 0$, $\kappa_1 \geq \kappa_N$, $\kappa_1 \geq 1$ and $\kappa_N > 0$, but we can prove the other cases in the same way. We define the exponent $\tilde{\kappa}$ as

$$h_{1\mathrm{op}} - h_{N\mathrm{op}} = O(\beta^{-\tilde{\kappa}}) \tag{13}$$

in the limit $\beta \to 0$ and consider the case

$$\tilde{\kappa} > 0 \tag{14}$$

in the following. We discuss the case $\tilde{\kappa} \leq 0$ in Appendix A. Here, in order to obtain the inequality (9), we prove the following; F_1 and F_2 are of order of

$$O(\beta^{\kappa_1+\kappa_N+\kappa}) \text{ and } O(\beta^{\kappa_1+\kappa_N+\kappa'}), \tag{15}$$

or higher, respectively, where

$$\kappa = \min(\kappa_N, 1) > 0 \text{ and } \kappa' = \min(\kappa_N, \tilde{\kappa}, 1) > 0. \tag{16}$$

On the other hand, $p_{\uparrow\uparrow}p_{\downarrow\downarrow}$ and $p_{\uparrow\downarrow}p_{\downarrow\uparrow}$ are both of order of

$$O(\beta^{2\kappa_1+2\kappa_N}), \tag{17}$$

or lower. Then, the inequality (9) is satisfied below a certain value of β.

In order to prove (15) and (17), we separate the total Hamiltonian as follows:

$$H_{\mathrm{tot}} = H_1 + H_{\mathrm{couple}} + H_{\mathrm{media}} + H_N, \tag{18}$$

where

$$\begin{aligned} H_1 &= h_{1\mathrm{op}}\sigma_1^z, \quad H_N = h_{N\mathrm{op}}\sigma_N^z, \\ H_{\mathrm{couple}} &= J^x\sigma_1^x\sigma_2^x + J^y\sigma_1^y\sigma_2^y + J^z\sigma_1^z\sigma_2^z + J^x\sigma_{N-1}^x\sigma_N^x + J^y\sigma_{N-1}^y\sigma_N^y + J^z\sigma_{N-1}^z\sigma_N^z, \\ H_{\mathrm{media}} &= \sum_{i=2}^{N-2}(J_i^x\sigma_i^x\sigma_{i+1}^x + J_i^y\sigma_i^y\sigma_{i+1}^y + J_i^z\sigma_i^z\sigma_{i+1}^z) + \sum_{i=2}^{N-1} h_i^z\sigma_i^z. \end{aligned} \tag{19}$$

Now, we consider the term $H_{\rm couple}$, which couples the focused spins and the media spins, as perturbation. The unperturbed density matrix $\rho^{(0)}_{\rm tot}$ is given by

$$\rho^{(0)}_{\rm tot} = e^{-\beta H_1 - \beta H_N} e^{-\beta H_{\rm media}} \tag{20}$$

because H_1, $H_{\rm media}$ and H_N commute with each other. Because the external fields are applied in the z direction, the eigenstates of $H_1 + H_N$ are given by $\{|{\uparrow_1}{\uparrow_N}\rangle, |{\uparrow_1}{\downarrow_N}\rangle, |{\downarrow_1}{\uparrow_N}\rangle, |{\downarrow_1}{\downarrow_N}\rangle\}$ with the corresponding eigenvalues $\{-h_{1\rm op} - h_{N\rm op}, -h_{1\rm op} + h_{N\rm op}, h_{1\rm op} - h_{N\rm op}, h_{1\rm op} + h_{N\rm op}\}$; we denote these eigenvalues as $\{E^{\uparrow\uparrow}_{1N}, E^{\uparrow\downarrow}_{1N}, E^{\downarrow\uparrow}_{1N}, E^{\downarrow\downarrow}_{1N}\}$. We also give the eigenstates of $H_{\rm media}$ as

$$|\psi^n_{\rm media}\rangle = s_n|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_n\rangle|{\uparrow_{N-1}}\rangle + t_n|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_n\rangle|{\downarrow_{N-1}}\rangle + u_n|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_n\rangle|{\uparrow_{N-1}}\rangle + w_n|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_n\rangle|{\downarrow_{N-1}}\rangle, \tag{21}$$

for $n = 1, 2, \cdots, 2^{N-2}$, where $\{|\tilde{\psi}^{\uparrow\uparrow}_n\rangle, |\tilde{\psi}^{\uparrow\downarrow}_n\rangle, |\tilde{\psi}^{\downarrow\uparrow}_n\rangle, |\tilde{\psi}^{\downarrow\downarrow}_n\rangle\}$ are the states of the spins from 3 to $N-2$. Because the total Hamiltonian $H_{\rm tot}$ is a real matrix, the parameters $\{s_n, t_n, u_n, w_n\}$ are real numbers. We define the corresponding eigenvalues of $H_{\rm media}$ as $\{E^n_{\rm media}\}$. Then, the unperturbed eigenstates are given by

$$\{|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi^n_{\rm media}\rangle, |{\uparrow_1}{\downarrow_N}\rangle \otimes |\psi^n_{\rm media}\rangle, |{\downarrow_1}{\uparrow_N}\rangle \otimes |\psi^n_{\rm media}\rangle, |{\downarrow_1}{\downarrow_N}\rangle \otimes |\psi^n_{\rm media}\rangle\}. \tag{22}$$

We define the perturbed eigenstates corresponding to each of (22) as $\{|\psi^{n,\uparrow\uparrow}_{\rm tot}\rangle, |\psi^{n,\uparrow\downarrow}_{\rm tot}\rangle, |\psi^{n,\downarrow\uparrow}_{\rm tot}\rangle, |\psi^{n,\downarrow\downarrow}_{\rm tot}\rangle\}$, respectively. We express them as

$$|\psi^{n,\xi}_{\rm tot}\rangle = |{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi^{n,\xi}_{{\rm media},\uparrow\uparrow}\rangle + |{\uparrow_1}{\downarrow_N}\rangle \otimes |\psi^{n,\xi}_{{\rm media},\uparrow\downarrow}\rangle + |{\downarrow_1}{\uparrow_N}\rangle \otimes |\psi^{n,\xi}_{{\rm media},\downarrow\uparrow}\rangle + |{\downarrow_1}{\downarrow_N}\rangle \otimes |\psi^{n,\xi}_{{\rm media},\downarrow\downarrow}\rangle, \tag{23}$$

for $n = 1, 2, \cdots, 2^{N-2}$ and $\xi = \uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$, where $|\psi^{n,\xi}_{{\rm media},\uparrow\uparrow}\rangle$, $|\psi^{n,\xi}_{{\rm media},\uparrow\downarrow}\rangle$, $|\psi^{n,\xi}_{{\rm media},\downarrow\uparrow}\rangle$ and $|\psi^{n,\xi}_{{\rm media},\downarrow\downarrow}\rangle$ are the states of the spins from 2 to $N-1$ and may be not normalized. We also define the corresponding perturbed eigenvalues as $\{E^{n,\uparrow\uparrow}_{\rm tot}, E^{n,\uparrow\downarrow}_{\rm tot}, E^{n,\downarrow\uparrow}_{\rm tot}, E^{n,\downarrow\downarrow}_{\rm tot}\}$, which we express as

$$E^{n,\xi}_{\rm tot} = E^{\xi}_{1N} + E^n_{\rm media} + \delta E^{n,\xi}_{\rm tot} \tag{24}$$

for $n = 1, 2, \cdots, 2^{N-2}$ and $\xi = \uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$. Note that $\{E^n_{\rm media}\}$ and $\{\delta E^{n,\xi}_{\rm tot}\}$ are of order β^0 because $H_{\rm media}$ and $H_{\rm couple}$ do not depend on the temperature β. Then, we have

$$\frac{e^{-\beta E^{n,\uparrow\uparrow}_{\rm tot}}}{Z_{\rm tot}} = \frac{e^{\beta(h_{1\rm op} + h_{N\rm op}) - \beta(E^n_{\rm media} + \delta E^{n,\uparrow\uparrow}_{\rm tot})}}{Z_{\rm tot}} = O(\beta^0) \tag{25}$$

in the limit $\beta \to 0$ in the case $h_{1\rm op}, h_{N\rm op} > 0$, where $Z_{\rm tot}$ is the partition function of the total Hamiltonian.

Now, we show the outline of the proof. First, we can calculate the elements $\{F_1, F_2, p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$ from (23) and (24) as

$$F_1 = \frac{1}{Z_{\rm tot}} \sum_{n=1}^{2^{N-2}} \sum_{\xi=\uparrow\uparrow,\uparrow\downarrow,\downarrow\uparrow,\downarrow\downarrow} e^{-\beta E^{n,\xi}_{\rm tot}} \langle \psi^{n,\xi}_{{\rm media},\uparrow\downarrow} | \psi^{n,\xi}_{{\rm media},\downarrow\uparrow} \rangle, \tag{26}$$

and

$$p_{\uparrow\uparrow} = \frac{1}{Z_{\rm tot}} \sum_{n=1}^{2^{N-2}} \sum_{\xi=\uparrow\uparrow,\uparrow\downarrow,\downarrow\uparrow,\downarrow\downarrow} e^{-\beta E^{n,\xi}_{\rm tot}} \langle \psi^{n,\xi}_{{\rm media},\uparrow\uparrow} | \psi^{n,\xi}_{{\rm media},\uparrow\uparrow} \rangle, \tag{27}$$

for example. Second, we estimate the leading orders of $\{F_1, F_2\}$ and $\{p_{\uparrow\downarrow}p_{\downarrow\uparrow}, p_{\uparrow\uparrow}p_{\downarrow\downarrow}\}$. From the perturbation theory, we can obtain the approximate forms of $\{|\psi^{n,\xi}_{\rm tot}\rangle\}$ and expand $\{F_1, F_2\}$ and $\{p_{\uparrow\downarrow}p_{\downarrow\uparrow}, p_{\uparrow\uparrow}p_{\downarrow\downarrow}\}$ with respect to β. The elements $\{F_1, F_2\}$ are additive with respect to the indices n and ξ; we define each contribution to the elements $\{F_1, F_2\}$ as

$$\begin{aligned} F^{n,\xi}_1 &= \langle \psi^{n,\xi}_{{\rm media},\uparrow\downarrow} | \psi^{n,\xi}_{{\rm media},\downarrow\uparrow} \rangle, \\ F^{n,\xi}_2 &= \langle \psi^{n,\xi}_{{\rm media},\uparrow\uparrow} | \psi^{n,\xi}_{{\rm media},\downarrow\downarrow} \rangle. \end{aligned} \tag{28}$$

Then, the elements $\{F_1, F_2\}$ are given by

$$\begin{aligned} F_1 &= \frac{1}{Z_{\mathrm{tot}}} \sum_{n=1}^{2^{N-2}} \sum_{\xi=\uparrow\uparrow,\uparrow\downarrow,\downarrow\uparrow,\downarrow\downarrow} e^{-\beta E_{\mathrm{tot}}^{n,\xi}} F_1^{n,\xi}, \\ F_2 &= \frac{1}{Z_{\mathrm{tot}}} \sum_{n=1}^{2^{N-2}} \sum_{\xi=\uparrow\uparrow,\uparrow\downarrow,\downarrow\uparrow,\downarrow\downarrow} e^{-\beta E_{\mathrm{tot}}^{n,\xi}} F_2^{n,\xi}, \end{aligned} \tag{29}$$

respectively. The elements $\{p_{\uparrow\downarrow}p_{\downarrow\uparrow}, p_{\uparrow\uparrow}p_{\downarrow\downarrow}\}$, on the other hand, are given by double summations as

$$\begin{aligned} p_{\uparrow\downarrow}p_{\downarrow\uparrow} &= \frac{1}{Z_{\mathrm{tot}}^2} \sum_{n,n'=1}^{2^{N-2}} \sum_{\xi,\xi'=\uparrow\uparrow,\uparrow\downarrow,\downarrow\uparrow,\downarrow\downarrow} e^{-\beta E_{\mathrm{tot}}^{n,\xi}} \langle \psi_{\mathrm{media},\uparrow\downarrow}^{n,\xi} | \psi_{\mathrm{media},\uparrow\downarrow}^{n,\xi} \rangle e^{-\beta E_{\mathrm{tot}}^{n',\xi'}} \langle \psi_{\mathrm{media},\downarrow\uparrow}^{n',\xi'} | \psi_{\mathrm{media},\downarrow\uparrow}^{n',\xi'} \rangle, \\ p_{\uparrow\uparrow}p_{\downarrow\downarrow} &= \frac{1}{Z_{\mathrm{tot}}^2} \sum_{n,n'=1}^{2^{N-2}} \sum_{\xi,\xi'=\uparrow\uparrow,\uparrow\downarrow,\downarrow\uparrow,\downarrow\downarrow} e^{-\beta E_{\mathrm{tot}}^{n,\xi}} \langle \psi_{\mathrm{media},\uparrow\uparrow}^{n,\xi} | \psi_{\mathrm{media},\uparrow\uparrow}^{n,\xi} \rangle e^{-\beta E_{\mathrm{tot}}^{n',\xi'}} \langle \psi_{\mathrm{media},\downarrow\downarrow}^{n',\xi'} | \psi_{\mathrm{media},\downarrow\downarrow}^{n',\xi'} \rangle. \end{aligned} \tag{30}$$

Because the summands of (30) are all positive, we can obtain the following inequalities for $\{p_{\uparrow\downarrow}p_{\downarrow\uparrow}, p_{\uparrow\uparrow}p_{\downarrow\downarrow}\}$:

$$\begin{aligned} p_{\uparrow\downarrow}p_{\downarrow\uparrow} \geq & \frac{1}{Z_{\mathrm{tot}}^2} \sum_{n=1}^{2^{N-2}} e^{-\beta E_{\mathrm{tot}}^{n,\uparrow\uparrow}} \langle \psi_{\mathrm{media},\uparrow\downarrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\uparrow\downarrow}^{n,\uparrow\uparrow} \rangle e^{-\beta E_{\mathrm{tot}}^{n,\uparrow\uparrow}} \langle \psi_{\mathrm{media},\downarrow\uparrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\downarrow\uparrow}^{n,\uparrow\uparrow} \rangle \equiv P_{\uparrow\downarrow,\downarrow\uparrow}^{\uparrow\uparrow}, \\ p_{\uparrow\uparrow}p_{\downarrow\downarrow} \geq & \frac{1}{Z_{\mathrm{tot}}^2} \sum_{n=1}^{2^{N-2}} e^{-\beta E_{\mathrm{tot}}^{n,\uparrow\uparrow}} \langle \psi_{\mathrm{media},\uparrow\uparrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\uparrow\uparrow}^{n,\uparrow\uparrow} \rangle e^{-\beta E_{\mathrm{tot}}^{n,\uparrow\uparrow}} \langle \psi_{\mathrm{media},\downarrow\downarrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\downarrow\downarrow}^{n,\uparrow\uparrow} \rangle \equiv P_{\uparrow\uparrow,\downarrow\downarrow}^{\uparrow\uparrow}, \end{aligned} \tag{31}$$

where we pick up only the terms of $n = n'$ and $\xi = \xi' = \uparrow\uparrow$ from the summations in (30). The elements $P_{\uparrow\downarrow,\downarrow\uparrow}^{\uparrow\uparrow}$ and $P_{\uparrow\uparrow,\downarrow\downarrow}^{\uparrow\uparrow}$ are additive with respect to n; we define each contribution to the elements $\{P_{\uparrow\downarrow,\downarrow\uparrow}^{\uparrow\uparrow}, P_{\uparrow\uparrow,\downarrow\downarrow}^{\uparrow\uparrow}\}$ as

$$\begin{aligned} P_{\uparrow\downarrow,\downarrow\uparrow}^{n,\uparrow\uparrow} &= \langle \psi_{\mathrm{media},\uparrow\downarrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\uparrow\downarrow}^{n,\uparrow\uparrow} \rangle \langle \psi_{\mathrm{media},\downarrow\uparrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\downarrow\uparrow}^{n,\uparrow\uparrow} \rangle, \\ P_{\uparrow\uparrow,\downarrow\downarrow}^{n,\uparrow\uparrow} &= \langle \psi_{\mathrm{media},\uparrow\uparrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\uparrow\uparrow}^{n,\uparrow\uparrow} \rangle \langle \psi_{\mathrm{media},\downarrow\downarrow}^{n,\uparrow\uparrow} | \psi_{\mathrm{media},\downarrow\downarrow}^{n,\uparrow\uparrow} \rangle. \end{aligned} \tag{32}$$

Then, the elements $\{P_{\uparrow\downarrow,\downarrow\uparrow}^{\uparrow\uparrow}, P_{\uparrow\uparrow,\downarrow\downarrow}^{\uparrow\uparrow}\}$ are given by

$$\begin{aligned} P_{\uparrow\downarrow,\downarrow\uparrow}^{\uparrow\uparrow} &= \frac{1}{Z_{\mathrm{tot}}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\mathrm{tot}}^{n,\uparrow\uparrow}} P_{\uparrow\downarrow,\downarrow\uparrow}^{n,\uparrow\uparrow}, \\ P_{\uparrow\uparrow,\downarrow\downarrow}^{\uparrow\uparrow} &= \frac{1}{Z_{\mathrm{tot}}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\mathrm{tot}}^{n,\uparrow\uparrow}} P_{\uparrow\uparrow,\downarrow\downarrow}^{n,\uparrow\uparrow}, \end{aligned} \tag{33}$$

respectively. In this way, we calculate each contribution to the elements $\{F_1, F_2\}$ and $\{P_{\uparrow\downarrow,\downarrow\uparrow}^{\uparrow\uparrow}, P_{\uparrow\uparrow,\downarrow\downarrow}^{\uparrow\uparrow}\}$ separately.

In accordance with the above outline, we first calculate $|\psi_{\mathrm{tot}}^{n_0,\uparrow\uparrow}\rangle$, namely the perturbed state of $|\uparrow_1\uparrow_N\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle$. In order to simplify the calculation, we rewrite the perturbation Hamiltonian H_{couple} as follows:

$$\begin{aligned} H_{\mathrm{couple}} = & \frac{1}{2} J \left[(1+\gamma)\sigma_1^x \sigma_2^x + (1-\gamma)\sigma_1^y \sigma_2^y \right] + J^z \sigma_1^z \sigma_2^z \\ & + \frac{1}{2} J \left[(1+\gamma)\sigma_{N-1}^x \sigma_N^x + (1-\gamma)\sigma_{N-1}^y \sigma_N^y \right] + J^z \sigma_{N-1}^z \sigma_N^z \\ = & J \left[\sigma_1^+ \sigma_2^- + \sigma_1^- \sigma_2^+ + \gamma(\sigma_1^+ \sigma_2^+ + \sigma_1^- \sigma_2^-) \right] + J^z \sigma_1^z \sigma_2^z \\ & + J \left[\sigma_{N-1}^+ \sigma_N^- + \sigma_{N-1}^- \sigma_N^+ + \gamma(\sigma_{N-1}^+ \sigma_N^+ + \sigma_{N-1}^- \sigma_N^-) \right] + J^z \sigma_{N-1}^z \sigma_N^z, \end{aligned} \tag{34}$$

where

$$J = J^x + J^y, \quad \gamma = \frac{J^x - J^y}{J^x + J^y}. \tag{35}$$

From the calculation in Ref. [20], the leading terms of the elements $\{|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\xi}\rangle\}$ for $\xi = \uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$ in Eq. (23) are given by

$$|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\uparrow\uparrow}\rangle = |\psi^{n_0}_{\mathrm{media}}\rangle, \tag{36}$$

$$\begin{aligned}|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\uparrow\downarrow}\rangle =& \frac{J}{-2h_{N\mathrm{op}}}\Big(\gamma s_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + t_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle \\ &+ \gamma u_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + w_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + O(\beta^{\kappa_N})\Big),\end{aligned} \tag{37}$$

$$\begin{aligned}|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\downarrow\uparrow}\rangle =& \frac{J}{-2h_{1\mathrm{op}}}\Big(\gamma s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle \\ &+ u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + O(\beta^{\kappa_1})\Big)\end{aligned} \tag{38}$$

and

$$\begin{aligned}|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\downarrow\downarrow}\rangle =& \frac{J^2}{4h_{1\mathrm{op}}h_{N\mathrm{op}}}\Big(\gamma^2 s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle \\ &+ \gamma u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + O(\beta^{\kappa_N})\Big).\end{aligned} \tag{39}$$

Now, we calculate the contributions of $|\psi^{n_0,\uparrow\uparrow}_{\mathrm{tot}}\rangle$ to the elements $\{F_1, F_2\}$ and $\{P^{\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}, P^{\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}\}$, which are defined as $\{F^{n_0,\uparrow\uparrow}_1, F^{n_0,\uparrow\uparrow}_2\}$ and $\{P^{n_0,\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}, P^{n_0,\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}\}$. From Eqs. (36)–(39), we have the elements $\{F^{n_0,\uparrow\uparrow}_1, F^{n_0,\uparrow\uparrow}_2\}$ as

$$\begin{aligned}F^{n_0,\uparrow\uparrow}_1 =& \frac{J^2}{4h_{1\mathrm{op}}h_{N\mathrm{op}}}\Big(\gamma s_{n_0}w_{n_0}\langle\tilde{\psi}^{\downarrow\downarrow}_{n_0}|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle + u_{n_0}t_{n_0}\langle\tilde{\psi}^{\uparrow\downarrow}_{n_0}|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle \\ &+ \gamma^2 t_{n_0}u_{n_0}\langle\tilde{\psi}^{\downarrow\uparrow}_{n_0}|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle + \gamma w_{n_0}s_{n_0}\langle\tilde{\psi}^{\uparrow\uparrow}_{n_0}|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle + O(\beta^{\kappa_N})\Big)\end{aligned} \tag{40}$$

and

$$\begin{aligned}F^{n_0,\uparrow\uparrow}_2 =& \frac{J^2}{4h_{1\mathrm{op}}h_{N\mathrm{op}}}\Big(\gamma^2 w_{n_0}s_{n_0}\langle\tilde{\psi}^{\downarrow\downarrow}_{n_0}|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle + \gamma u_{n_0}t_{n_0}\langle\tilde{\psi}^{\downarrow\uparrow}_{n_0}|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle \\ &+ \gamma t_{n_0}u_{n_0}\langle\tilde{\psi}^{\uparrow\downarrow}_{n_0}|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle + s_{n_0}w_{n_0}\langle\tilde{\psi}^{\uparrow\uparrow}_{n_0}|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle + O(\beta^{\kappa_N})\Big)\end{aligned} \tag{41}$$

as well as the elements $\{P^{n_0,\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}, P^{n_0,\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}\}$ in the forms

$$\begin{aligned}P^{n_0,\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow} =& \frac{J^4}{16h^2_{1\mathrm{op}}h^2_{N\mathrm{op}}}\Big(\gamma^2 s^2_{n_0} + t^2_{n_0} + \gamma^2 u^2_{n_0} + w^2_{n_0} + O\big(\beta^{\kappa_N}\big)\Big) \\ &\times\Big(\gamma^2 s^2_{n_0} + \gamma^2 t^2_{n_0} + u^2_{n_0} + w^2_{n_0} + O(\beta^{\kappa_N})\Big) \\ \geq& \frac{J^4}{16h^2_{1\mathrm{op}}h^2_{N\mathrm{op}}}(w^4_{n_0} + \beta^{\kappa_N}\delta P^{n_0,\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow})\end{aligned} \tag{42}$$

and

$$\begin{aligned}P^{n_0,\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow} =& \frac{J^4}{16h^2_{1\mathrm{op}}h^2_{N\mathrm{op}}}\Big(\gamma^4 s^2_{n_0} + \gamma^2 t^2_{n_0} + \gamma^2 u^2_{n_0} + w^2_{n_0} + O(\beta^{\kappa_N})\Big) \\ \geq& \frac{J^4}{16h^2_{1\mathrm{op}}h^2_{N\mathrm{op}}}(w^2_{n_0} + \beta^{\kappa_N}\delta P^{n_0,\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow})\end{aligned} \tag{43}$$

in the limit $\beta \to 0$, where we define the real numbers of order β^0 as $\delta P^{n_0,\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}$ and $\delta P^{n_0,\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}$ so as to satisfy the above inequalities.

Then, we sum the elements $\{F^{n,\uparrow\uparrow}_1, F^{n,\uparrow\uparrow}_2\}$ and $\{P^{n,\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}, P^{n,\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}\}$ with the Boltzmann weight $e^{-\beta E^{n,\uparrow\uparrow}_{\mathrm{tot}}}$ over the label n. First, we calculate the summation of each of $\{F^{n,\uparrow\uparrow}_1, F^{n,\uparrow\uparrow}_2\}$. Because the spins 2 and $(N-1)$ are separated by

$(N-4)$ spins, the correlation between the spins 2 and $(N-1)$ are generated by the $(N-3)$th-order perturbation of $H_{\rm media}$. Therefore, we obtain

$$\begin{aligned}&\langle\sigma_2^x\sigma_{N-1}^x\rangle_0=O(\beta^{\alpha_1}),\\&\langle\sigma_2^y\sigma_{N-1}^y\rangle_0=O(\beta^{\alpha_2}),\end{aligned}\tag{44}$$

where $\langle\cdots\rangle_0$ denotes the thermal average with respect to $\rho_{\rm tot}^{(0)}$ in (20) and $\alpha_1\geq N-3$, $\alpha_2\geq N-3$. Since we are considering the case $N\geq 4$, we have $\alpha_1\geq 1$ and $\alpha_2\geq 1$. This is the key to the fact that the entanglement can vanish for $N\geq 4$. From the equations

$$\begin{aligned}&\frac{\langle\sigma_2^x\sigma_{N-1}^x+\sigma_2^y\sigma_{N-1}^y\rangle_0}{4}\\=&\frac{\langle\sigma_2^+\sigma_{N-1}^-+\sigma_2^-\sigma_{N-1}^+\rangle_0}{2}\\=&{\rm tr}\left(e^{-\beta H_{\rm media}}\frac{|{\downarrow_2}{\uparrow_{N-1}}\rangle\langle{\uparrow_2}{\downarrow_{N-1}}|+|{\uparrow_2}{\downarrow_{N-1}}\rangle\langle{\downarrow_2}{\uparrow_{N-1}}|}{2}\right)\\=&{\rm tr}\Big(e^{-\beta H_{\rm media}}|{\downarrow_2}{\uparrow_{N-1}}\rangle\langle{\uparrow_2}{\downarrow_{N-1}}|\Big)\\=&\frac{1}{Z_{\rm media}}\sum_{n=1}^{2^{N-2}}e^{-\beta E^n_{\rm media}}\big\langle{\uparrow_2}{\downarrow_{N-1}}\,\big|\psi^n_{\rm media}\big\rangle\big\langle\psi^n_{\rm media}\big|{\downarrow_2}{\uparrow_{N-1}}\big\rangle\end{aligned}\tag{45}$$

and

$$\begin{aligned}&\frac{\langle\sigma_2^x\sigma_{N-1}^x-\sigma_2^y\sigma_{N-1}^y\rangle_0}{4}\\=&\frac{\langle\sigma_2^+\sigma_{N-1}^++\sigma_2^-\sigma_{N-1}^-\rangle_0}{2}\\=&{\rm tr}\left(e^{-\beta H_{\rm media}}\frac{|{\downarrow_2}{\downarrow_{N-1}}\rangle\langle{\uparrow_2}{\uparrow_{N-1}}|+|{\uparrow_2}{\uparrow_{N-1}}\rangle\langle{\downarrow_2}{\downarrow_{N-1}}|}{2}\right)\\=&{\rm tr}\Big(e^{-\beta H_{\rm media}}|{\uparrow_2}{\uparrow_{N-1}}\rangle\langle{\downarrow_2}{\downarrow_{N-1}}|\Big)\\=&\frac{1}{Z_{\rm media}}\sum_{n=1}^{2^{N-2}}e^{-\beta E^n_{\rm media}}\big\langle{\downarrow_2}{\downarrow_{N-1}}\,\big|\psi^n_{\rm media}\big\rangle\big\langle\psi^n_{\rm media}\big|{\uparrow_2}{\uparrow_{N-1}}\big\rangle,\end{aligned}\tag{46}$$

where $Z_{\rm media}\equiv{\rm tr}(e^{-\beta H_{\rm media}})$, $\sigma^+\equiv(\sigma^x+i\sigma^y)/2$ and $\sigma^-\equiv(\sigma^x-i\sigma^y)/2$, we also have

$$\sum_{n=1}^{2^{N-2}}e^{-\beta E^n_{\rm media}}t_nu_n\langle\tilde{\psi}_n^{\uparrow\downarrow}|\tilde{\psi}_n^{\downarrow\uparrow}\rangle=O(\beta^\alpha)\tag{47}$$

and

$$\sum_{n=1}^{2^{N-2}}e^{-\beta E^n_{\rm media}}s_nw_n\langle\tilde{\psi}_n^{\downarrow\downarrow}|\tilde{\psi}_n^{\uparrow\uparrow}\rangle=O(\beta^\alpha),\tag{48}$$

where $\alpha=\min(\alpha_1,\alpha_2)$. Moreover, because $\{\delta E_{\rm tot}^{n,\xi}\}$ in Eq. (24) are of order β^0, we have

$$e^{-\beta E_{\rm tot}^{n,\xi}}=e^{-\beta(E_{1N}^\xi+E^n_{\rm media})}\Big(1-\beta\delta E_{\rm tot}^{n,\xi}+O(\beta^2)\Big),\tag{49}$$

for $n = 1, 2, \cdots, 2^{N-2}$ and $\xi = \uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$. Then, we obtain from Eqs. (40), (41), and (47)–(49),

$$\begin{aligned}
&\sum_{n=1}^{2^{N-2}} e^{-\beta E_{\rm tot}^{n,\uparrow\uparrow}} F_1^{n,\uparrow\uparrow} \\
&= \sum_{n=1}^{2^{N-2}} e^{-\beta(E_{1N}^{\uparrow\uparrow}+E_{\rm media}^{n})} F_1^{n,\uparrow\uparrow} + \sum_{n=1}^{2^{N-2}} e^{-\beta(E_{1N}^{\uparrow\uparrow}+E_{\rm media}^{n})} \big(-\beta\delta E_{\rm tot}^{n,\uparrow\uparrow} + O(\beta^2)\big) F_1^{n,\uparrow\uparrow} \\
&= e^{-\beta E_{1N}^{\uparrow\uparrow}} \frac{J^2}{4h_{1\rm op}h_{N\rm op}} \Big(O(\beta^{\alpha}) + O(\beta^{\kappa_N})\Big) + e^{-\beta E_{1N}^{\uparrow\uparrow}} \frac{J^2}{4h_{1\rm op}h_{N\rm op}} O(\beta) \\
&= e^{-\beta E_{1N}^{\uparrow\uparrow}} \times O(\beta^{\kappa_1+\kappa_N+\kappa})
\end{aligned} \tag{50}$$

and

$$\sum_{n=1}^{2^{N-2}} e^{-\beta E_{\rm tot}^{n,\uparrow\uparrow}} F_2^{n,\uparrow\uparrow} = e^{-\beta E_{1N}^{\uparrow\uparrow}} \frac{J^2}{4h_{1\rm op}h_{N\rm op}} \Big(O(\beta^{\alpha}) + O(\beta^{\kappa_N}) + O(\beta)\Big) = e^{-\beta E_{1N}^{\uparrow\uparrow}} \times O(\beta^{\kappa_1+\kappa_N+\kappa}), \tag{51}$$

where $\alpha = \min(\alpha_1, \alpha_2) \geq 1$ and κ is defined in (16).

We can similarly calculate the contributions of the other states $\{|\psi_{\rm tot}^{n,\xi}\rangle\}$ ($\xi = \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$). From the calculation in Ref. [20], the leading terms of the elements $\{|\psi_{{\rm media},\xi}^{n_0,\uparrow\downarrow}\rangle\}$ for $\xi = \uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$ in Eq. (23) can be obtained, and we have the contributions of the states $\{|\psi_{\rm tot}^{n,\uparrow\downarrow}\rangle\}$ to F_1 and F_2 as

$$\begin{aligned}
\sum_{n=1}^{2^{N-2}} e^{-\beta E_{\rm tot}^{n,\uparrow\downarrow}} F_1^{n,\uparrow\downarrow} &= e^{-\beta E_{1N}^{\uparrow\downarrow}} \frac{J^2}{4h_{1\rm op}h_{N\rm op}} \Big(O(\beta^{\alpha}) + O(\beta^{\kappa_N}) + O(\beta^{\tilde{\kappa}}) + O(\beta)\Big) = e^{-\beta E_{1N}^{\uparrow\downarrow}} \times O(\beta^{\kappa_1+\kappa_N+\kappa'}), \\
\sum_{n=1}^{2^{N-2}} e^{-\beta E_{\rm tot}^{n,\uparrow\downarrow}} F_2^{n,\uparrow\downarrow} &= e^{-\beta E_{1N}^{\uparrow\downarrow}} \frac{J^2}{4h_{1\rm op}h_{N\rm op}} \Big(O(\beta^{\alpha}) + O(\beta^{\kappa_N}) + O(\beta)\Big) = e^{-\beta E_{1N}^{\uparrow\downarrow}} \times O(\beta^{\kappa_1+\kappa_N+\kappa}),
\end{aligned} \tag{52}$$

where we utilized Eqs. (47)–(49) and κ' is defined in (16). From the inequality

$$\frac{e^{-\beta E_{1N}^{\xi}}}{Z_{\rm tot}} < 1 \tag{53}$$

for $\xi = \uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$, we finally obtain (15) by substituting Eqs. (50)–(52) into Eq. (29).

Second, we calculate the summation of each of $\{P_{\uparrow\downarrow,\downarrow\uparrow}^{n,\uparrow\uparrow}, P_{\uparrow\uparrow,\downarrow\downarrow}^{n,\uparrow\uparrow}\}$. The parameter w_n in Eq. (21) cannot vanish for all n. Therefore, we have

$$\begin{aligned}
\frac{1}{Z_{\rm tot}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\rm tot}^{n,\uparrow\uparrow}} w_n^4 &= O(\beta^0), \\
\frac{1}{Z_{\rm tot}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\rm tot}^{n,\uparrow\uparrow}} w_n^2 &= O(\beta^0),
\end{aligned} \tag{54}$$

where we utilized Eq. (25). Hence we obtain

$$\begin{aligned}
P_{\uparrow\downarrow,\downarrow\uparrow}^{\uparrow\uparrow} &= \frac{1}{Z_{\rm tot}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\rm tot}^{n,\uparrow\uparrow}} P_{\uparrow\downarrow,\downarrow\uparrow}^{n,\uparrow\uparrow} \\
&\geq \frac{J^4}{16h_{1\rm op}^2 h_{N\rm op}^2} \frac{1}{Z_{\rm tot}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\rm tot}^{n,\uparrow\uparrow}} (w_n^4 + \beta^{\kappa_N} \delta P_{\uparrow\downarrow,\downarrow\uparrow}^{n,\uparrow\uparrow}) = O(\beta^{2\kappa_1+2\kappa_N})
\end{aligned} \tag{55}$$

and

$$\begin{aligned}
P_{\uparrow\uparrow,\downarrow\downarrow}^{\uparrow\uparrow} &= \frac{1}{Z_{\rm tot}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\rm tot}^{n,\uparrow\uparrow}} P_{\uparrow\uparrow,\downarrow\downarrow}^{n,\uparrow\uparrow} \\
&\geq \frac{J^4}{16h_{1\rm op}^2 h_{N\rm op}^2} \frac{1}{Z_{\rm tot}^2} \sum_{n=1}^{2^{N-2}} e^{-2\beta E_{\rm tot}^{n,\uparrow\uparrow}} (w_n^2 + \beta^{\kappa_N} \delta P_{\uparrow\uparrow,\downarrow\downarrow}^{n,\uparrow\uparrow}) = O(\beta^{2\kappa_1+2\kappa_N}).
\end{aligned} \tag{56}$$

Because $p_{\uparrow\downarrow}p_{\downarrow\uparrow} \geq P^{\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}$ and $p_{\uparrow\uparrow}p_{\downarrow\downarrow} \geq P^{\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}$, we obtain (17). Thus, the inequality (9) is satisfied below a certain value of β in the case (b).

Case (c). To simplify the problem, we consider the case of $h_1 > 0$, $\kappa_1 \geq 1$ and $\kappa_N \leq 0$, but we can prove the other cases in the same way. In this case, we prove the following; F_1^2 and F_2^2 are both of order of

$$O(\beta^{2\kappa_1+2}), \tag{57}$$

or higher. On the other hand, $p_{\uparrow\uparrow}p_{\downarrow\downarrow}$ and $p_{\uparrow\downarrow}p_{\downarrow\uparrow}$ are both of order of

$$O(\beta^{2\kappa_1}), \tag{58}$$

or lower. Then, the inequality (9) is satisfied below a certain value of β.

In a similar manner to the case (b), we separate the total Hamiltonian as follows:

$$H_{\mathrm{tot}} = H_1 + H'_{\mathrm{couple}} + H'_{\mathrm{media}}, \tag{59}$$

where

$$\begin{aligned} &H_1 = h_{1\mathrm{op}}\sigma_1^z, \quad H'_{\mathrm{couple}} = J^x\sigma_1^x\sigma_2^x + J^y\sigma_1^y\sigma_2^y + J^z\sigma_1^z\sigma_2^z, \\ &H'_{\mathrm{media}} = \sum_{i=2}^{N-1}(J_i^x\sigma_i^x\sigma_{i+1}^x + J_i^y\sigma_i^y\sigma_{i+1}^y + J_i^z\sigma_i^z\sigma_{i+1}^z) + \sum_{i=2}^{N-1} h_i^z\sigma_i^z + h_{N\mathrm{op}}\sigma_N^z. \end{aligned} \tag{60}$$

Note that in the case (c), the norm of the Hamiltonian H'_{media} is of order β^0 because $\kappa_N < 0$. Here, we consider the interaction term H'_{couple}, which couples the spins 1 and the other spins, as perturbation. Then, the unperturbed density matrix $\rho_{\mathrm{tot}}^{'(0)}$ is expressed as

$$\rho_{\mathrm{tot}}^{'(0)} = e^{-\beta H_1}e^{-\beta H'_{\mathrm{media}}}. \tag{61}$$

The eigenstates of H_1 can be given by $\{|\uparrow_1\rangle, |\downarrow_1\rangle\}$ with the corresponding eigenvalues $\{-h_{1\mathrm{op}}, h_{1\mathrm{op}}\}$; we denote these eigenvalues as $\{E_1^{\uparrow}, E_1^{\downarrow}\}$. We also define the eigenstates of H'_{media} as

$$|\phi^n_{\mathrm{media}}\rangle = s'_n|\uparrow_2\rangle|\tilde{\phi}_n^{\uparrow\uparrow}\rangle|\uparrow_N\rangle + t'_n|\uparrow_2\rangle|\tilde{\phi}_n^{\uparrow\downarrow}\rangle|\downarrow_N\rangle + u'_n|\downarrow_2\rangle|\tilde{\phi}_n^{\downarrow\uparrow}\rangle|\uparrow_N\rangle + w'_n|\downarrow_2\rangle|\tilde{\phi}_n^{\downarrow\downarrow}\rangle|\downarrow_N\rangle \tag{62}$$

for $n = 1, 2, \cdots 2^{N-1}$, where $\{|\tilde{\phi}_n^{\uparrow\uparrow}\rangle, |\tilde{\phi}_n^{\uparrow\downarrow}\rangle, |\tilde{\phi}_n^{\downarrow\uparrow}\rangle, |\tilde{\phi}_n^{\downarrow\downarrow}\rangle\}$ are the states of the spins from 3 to $N-1$. Because the total Hamiltonian H_{tot} is a real matrix, the parameters $\{s'_n, t'_n, u'_n, w'_n\}$ are real numbers. We define the corresponding eigenvalues of H'_{media} as $\{E^{'n}_{\mathrm{media}}\}$. Then, the unperturbed eigenstates are given by $\{|\uparrow_1\rangle \otimes |\phi^n_{\mathrm{media}}\rangle\}$ and $\{|\downarrow_1\rangle \otimes |\phi^n_{\mathrm{media}}\rangle\}$. We define the corresponding perturbed eigenstates as $\{|\phi^{n,\uparrow}_{\mathrm{tot}}\rangle\}$ and $\{|\phi^{n,\downarrow}_{\mathrm{tot}}\rangle\}$ and the corresponding perturbed eigenvalues as $\{E^{n,\uparrow}_{\mathrm{tot}}, E^{n,\downarrow}_{\mathrm{tot}}\}$. We express them in the forms

$$|\phi^{n,\eta}_{\mathrm{tot}}\rangle = |\uparrow_1\rangle \otimes |\phi^{n,\eta}_{\mathrm{media},\uparrow}\rangle + |\downarrow_1\rangle \otimes |\phi^{n,\eta}_{\mathrm{media},\downarrow}\rangle \tag{63}$$

and

$$E^{'n,\eta}_{\mathrm{tot}} = E_1^{\eta} + E^{'n}_{\mathrm{media}} + \delta E^{'n,\eta}_{\mathrm{tot}} \tag{64}$$

for $n = 1, 2, \cdots 2^{N-1}$ and $\eta =\uparrow, \downarrow$, where $|\phi^{n,\eta}_{\mathrm{media},\uparrow}\rangle$ and $|\phi^{n,\eta}_{\mathrm{media},\downarrow}\rangle$ are the states of the spins from 2 to N and may not be normalized. Note that $\{E^n_{\mathrm{media}}\}$ and $\{\delta E^{'n,\xi}_{\mathrm{tot}}\}$ are of order β^0 because $||H'_{\mathrm{media}}||$ and $||H'_{\mathrm{couple}}||$ are of order β^0. Then, in the limit $\beta \to 0$, we have

$$\frac{e^{-\beta E^{'n,\uparrow}_{\mathrm{tot}}}}{Z_{\mathrm{tot}}} = \frac{e^{\beta h_{1\mathrm{op}} - \beta(E^{'n}_{\mathrm{media}} + \delta E^{'n,\uparrow}_{\mathrm{tot}})}}{Z_{\mathrm{tot}}} = O(\beta^0) \tag{65}$$

in the case $h_{1\mathrm{op}} > 0$.

Next, we calculate the elements $\{F_1, F_2\}$ and $\{p_{\uparrow\downarrow}p_{\downarrow\uparrow}, p_{\uparrow\uparrow}p_{\downarrow\downarrow}\}$. The calculation in [21] gives the state $|\phi^{n_0,\uparrow}_{\mathrm{tot}}\rangle$ in the form

$$\begin{aligned} |\phi^{n_0,\uparrow}_{\mathrm{tot}}\rangle = |\uparrow_1\rangle \otimes |\phi^{n_0}_{\mathrm{media}}\rangle + \frac{J}{-2h_{1\mathrm{op}}}|\downarrow_1\rangle \otimes \Big(&\gamma s'_{n_0}|\downarrow_2\rangle|\tilde{\phi}_{n_0}^{\uparrow\uparrow}\rangle|\uparrow_N\rangle + \gamma t'_{n_0}|\downarrow_2\rangle|\tilde{\phi}_{n_0}^{\uparrow\downarrow}\rangle|\downarrow_N\rangle \\ &+ u'_{n_0}|\uparrow_2\rangle|\tilde{\phi}_{n_0}^{\downarrow\uparrow}\rangle|\uparrow_N\rangle + w'_{n_0}|\uparrow_2\rangle|\tilde{\phi}_{n_0}^{\downarrow\downarrow}\rangle|\downarrow_N\rangle + O(\beta^{\kappa_1})\Big). \end{aligned} \tag{66}$$

Then, we can calculate the contribution of $|\phi_{\rm tot}^{n_0,\uparrow}\rangle$ to the elements $\{F_1, F_2\}$ and $\{p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$; we denote them as $\{F_1^{n_0,\uparrow}, F_2^{n_0,\uparrow}\}$ and $\{p_{\uparrow\uparrow}^{n_0,\uparrow}, p_{\uparrow\downarrow}^{n_0,\uparrow}, p_{\downarrow\uparrow}^{n_0,\uparrow}, p_{\downarrow\downarrow}^{n_0,\uparrow}\}$. First, the elements $\{F_1^{n_0,\uparrow}, F_2^{n_0,\uparrow}\}$ are given by

$$F_1^{n_0,\uparrow} = \frac{J}{-2h_{\rm 1op}}\Big(\gamma s'_{n_0} w'_{n_0} \langle\phi_{n_0}^{\downarrow\downarrow}|\phi_{n_0}^{\uparrow\uparrow}\rangle + u'_{n_0} t'_{n_0} \langle\phi_{n_0}^{\downarrow\uparrow}|\phi_{n_0}^{\uparrow\downarrow}\rangle + O(\beta^{\kappa_1})\Big) \tag{67}$$

and

$$F_2^{n_0,\uparrow} = \frac{J}{-2h_{\rm 1op}}\Big(\gamma t'_{n_0} u'_{n_0} \langle\phi_{n_0}^{\downarrow\uparrow}|\phi_{n_0}^{\uparrow\downarrow}\rangle + w'_{n_0} s'_{n_0} \langle\phi_{n_0}^{\uparrow\uparrow}|\phi_{n_0}^{\downarrow\downarrow}\rangle + O(\beta^{\kappa_1})\Big). \tag{68}$$

Second, the elements $\{p_{\uparrow\uparrow}^{n_0,\uparrow}, p_{\uparrow\downarrow}^{n_0,\uparrow}, p_{\downarrow\uparrow}^{n_0,\uparrow}, p_{\downarrow\downarrow}^{n_0,\uparrow}\}$ are given by

$$\begin{aligned} p_{\uparrow\uparrow}^{n_0,\uparrow} &= s'^2_{n_0} + u'^2_{n_0}, \\ p_{\uparrow\downarrow}^{n_0,\uparrow} &= t'^2_{n_0} + w'^2_{n_0}, \\ p_{\downarrow\uparrow}^{n_0,\uparrow} &= \frac{J^2}{4h_{\rm 1op}^2}\big(\gamma^2 s'^2_{n_0} + u'^2_{n_0} + O(\beta^{\kappa_1})\big), \\ p_{\downarrow\downarrow}^{n_0,\uparrow} &= \frac{J^2}{4h_{\rm 1op}^2}\big(\gamma^2 t'^2_{n_0} + w'^2_{n_0} + O(\beta^{\kappa_1})\big). \end{aligned} \tag{69}$$

Then, we sum $\{F_1^{n,\uparrow}, F_2^{n,\uparrow}\}$ and $\{p_{\uparrow\uparrow}^{n,\uparrow}, p_{\uparrow\downarrow}^{n,\uparrow}, p_{\downarrow\uparrow}^{n,\uparrow}, p_{\downarrow\downarrow}^{n,\uparrow}\}$ with the Boltzmann weight $e^{-\beta E_{\rm tot}^{'n,\uparrow}}$ over the label n. First, we calculate the summation of each of $\{F_1^{n,\uparrow}, F_2^{n,\uparrow}\}$. From the same discussion as in Eqs. (47) and (48) in the case (b), we have

$$\sum_{n=1}^{2^{N-1}} e^{-\beta E_{\rm media}^{'n}} t'_n u'_n \langle\tilde{\phi}_n^{\uparrow\downarrow}|\tilde{\phi}_n^{\downarrow\uparrow}\rangle = O(\beta^{\alpha'}) \tag{70}$$

and

$$\sum_{n=1}^{2^{N-1}} e^{-\beta E_{\rm media}^{'n}} s'_n w'_n \langle\tilde{\phi}_n^{\downarrow\downarrow}|\tilde{\phi}_n^{\uparrow\uparrow}\rangle = O(\beta^{\alpha'}), \tag{71}$$

where the exponent α' is defined as follows:

$$\begin{aligned} &\alpha' = \min(\alpha'_1, \alpha'_2), \\ &\langle\sigma_2^x\sigma_N^x\rangle = O(\beta^{\alpha'_1}), \quad \langle\sigma_2^y\sigma_N^y\rangle = O(\beta^{\alpha'_2}), \end{aligned} \tag{72}$$

where $\alpha'_1 \geq N-2$ and $\alpha'_2 \geq N-2$. Since we are considering the case $N \geq 4$, we have $\alpha'_1 \geq 2$ and $\alpha'_2 \geq 2$. On the other hand, because $\{\delta E_{\rm tot}^{'n,\eta}\}$ in Eq. (64) are of order β^0, we have

$$e^{-\beta E_{\rm tot}^{'n,\eta}} = e^{-\beta(E_1^\eta + E_{\rm media}^{'n})}\Big(1 - \beta\delta E_{\rm tot}^{'n,\eta} + O(\beta^2)\Big), \tag{73}$$

for $n = 1, 2, \cdots, 2^{N-1}$ and $\eta = \uparrow, \downarrow$. Then, we obtain from Eqs. (67), (68), and (70)–(73),

$$\begin{aligned} &\sum_{n=1}^{2^{N-1}} e^{-\beta E_{\rm tot}^{'n,\uparrow}} F_1^{n_0,\uparrow} \\ =&\sum_{n=1}^{2^{N-1}} e^{-\beta(E_1^\uparrow + E_{\rm media}^{'n})} F_1^{n_0,\uparrow} + \sum_{n=1}^{2^{N-1}} e^{-\beta(E_1^\uparrow + E_{\rm media}^{'n})}\Big(-\beta\delta E_{\rm tot}^{'n,\uparrow} + O(\beta^2)\Big) F_1^{n_0,\uparrow} \\ =&e^{-\beta E_1^\uparrow}\frac{J}{-2h_{\rm 1op}}\Big(O(\beta^{\alpha'}) + O(\beta^{\kappa_1})\Big) + e^{-\beta E_1^\uparrow}\frac{J}{-2h_{\rm 1op}} O(\beta) = e^{-\beta E_1^\uparrow} \times O(\beta^{\kappa_1+1}) \end{aligned} \tag{74}$$

and

$$\sum_{n=1}^{2^{N-1}} e^{-\beta E_{\rm tot}^{'n,\uparrow}} F_2^{n_0,\uparrow} = e^{-\beta E_1^\uparrow}\frac{J}{-2h_{\rm 1op}}\Big(O(\beta^{\alpha'}) + O(\beta^{\kappa_1}) + O(\beta)\Big) = e^{-\beta E_1^\uparrow} \times O(\beta^{\kappa_1+1}). \tag{75}$$

We similarly calculate the contributions of the other states $\{|\psi_{\mathrm{tot}}^{n,\downarrow}\rangle\}$; then, we finally arrive at (57),

$$
\begin{aligned}
F_1 &= \frac{1}{Z_{\mathrm{tot}}}\sum_{\eta=\uparrow,\downarrow} e^{-\beta E_1^\eta} \times O(\beta^{\kappa_1+1}) = O(\beta^{\kappa_1+1}),\\
F_2 &= \frac{1}{Z_{\mathrm{tot}}}\sum_{\eta=\uparrow,\downarrow} e^{-\beta E_1^\eta} \times O(\beta^{\kappa_1+1}) = O(\beta^{\kappa_1+1}),
\end{aligned}
\tag{76}
$$

where we utilized the inequality

$$
\frac{e^{-\beta E_1^\eta}}{Z_{\mathrm{tot}}} < 1 \tag{77}
$$

for $\eta =\uparrow,\downarrow$.

Second, we calculate the summation of each of $\{p_{\uparrow\uparrow}^{n,\uparrow}, p_{\uparrow\downarrow}^{n,\uparrow}, p_{\downarrow\uparrow}^{n,\uparrow}, p_{\downarrow\downarrow}^{n,\uparrow}\}$. From Eq. (69), we obtain

$$
\begin{aligned}
&\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\uparrow\uparrow}^{n,\uparrow} = e^{-\beta E_1^\uparrow} \times O(\beta^0),\\
&\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\uparrow\downarrow}^{n,\uparrow} = e^{-\beta E_1^\uparrow} \times O(\beta^0),\\
&\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\downarrow\uparrow}^{n,\uparrow} = e^{-\beta E_1^\uparrow} \times O(\beta^{2\kappa_1}),\\
&\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\downarrow\downarrow}^{n,\uparrow} = e^{-\beta E_1^\uparrow} \times O(\beta^{2\kappa_1}),
\end{aligned}
\tag{78}
$$

because $s_n'^2 + u_n'^2$, $t_n'^2 + w_n'^2$, $\gamma^2 s_n'^2 + u_n'^2$ and $\gamma^2 t_n'^2 + w_n'^2$ cannot vanish for all n. From Eq. (65), we have

$$
\begin{aligned}
p_{\uparrow\uparrow} &\geq \frac{1}{Z_{\mathrm{tot}}}\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\uparrow\uparrow}^{n,\uparrow} = O(\beta^0),\\
p_{\uparrow\downarrow} &\geq \frac{1}{Z_{\mathrm{tot}}}\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\uparrow\downarrow}^{n,\uparrow} = O(\beta^0),\\
p_{\downarrow\uparrow} &\geq \frac{1}{Z_{\mathrm{tot}}}\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\downarrow\uparrow}^{n,\uparrow} = O(\beta^{2\kappa_1}),\\
p_{\downarrow\downarrow} &\geq \frac{1}{Z_{\mathrm{tot}}}\sum_{n=1}^{2^{N-1}} e^{-\beta {E'}_{\mathrm{tot}}^{n,\uparrow}} p_{\downarrow\downarrow}^{n,\uparrow} = O(\beta^{2\kappa_1}).
\end{aligned}
\tag{79}
$$

We thereby obtain (57). We thus obtain (57) and (58) and hence the inequality (9) is satisfied below a certain value of β in the case (c).

Thus, we prove the inequality (9) in the cases (a), (b) and (c). This completes the proof of Theorem 1.

In this theorem, we discuss the case of $\{J_1^x, J_1^y, J_1^z\} = \{J_{N-1}^x, J_{N-1}^y, J_{N-1}^z\} = \{J^x, J^y, J^z\}$, but the case of $\{J_1^x, J_1^y, J_1^z\} \neq \{J_{N-1}^x, J_{N-1}^y, J_{N-1}^z\}$ can be also proved by extending this following proof; we only have to repeat the same calculation by changing H_{couple} in Eq. (19) into

$$
H_{\mathrm{couple}} = J_1^x\sigma_1^x\sigma_2^x + J_1^y\sigma_1^y\sigma_2^y + J_1^z\sigma_1^z\sigma_2^z + J_{N-1}^x\sigma_{N-1}^x\sigma_N^x + J_{N-1}^y\sigma_{N-1}^y\sigma_N^y + J_{N-1}^z\sigma_{N-1}^z\sigma_N^z. \tag{80}
$$

IV. ENTANGLEMENT MAXIMIZATION IN THREE-SPIN CHAINS

In the present section, we discuss the maximization problem in three-spin chains. A significant result of the present section is that the maximizing local fields are asymmetric to each other in a particular parameter region. This effect is



FIG. 2: Schematic picture of the three-spin chain. We define the spins 1 and 3 as the focused spins and the spin 2 as the media spin. The spins 1 and 3 are symmetric to each other and they indirectly interact with each other through the spin 2.

nontrivial; the positions of the spins 1 and 3 are geometrically symmetric to each other if $\{J_1^x, J_1^y, J_1^z\} = \{J_2^x, J_2^y, J_2^z\}$ and the entanglement is invariant with respect to the exchange of the two spins. Nevertheless, we obtain $|h_{1\mathrm{op}}| \neq |h_{3\mathrm{op}}|$ in a region. We show that the asymmetry can be mainly attributed to the behavior of the magnon. This effect indicates that the purity of the focused spins is not the only criterion for the enhancement of the bipartite entanglement. This is different from the case of the entanglement maximization in two-spin systems [14].

A. Numerical results

The Hamiltonian which we consider in the present section is

$$H_{\mathrm{tot}} = \sum_{i=1}^{2} (J^x \sigma_i^x \sigma_{i+1}^x + J^y \sigma_i^y \sigma_{i+1}^y + J^z \sigma_i^z \sigma_{i+1}^z) \\ + h_1^z \sigma_1^z + h^z_{\mathrm{media}} \sigma_2^z + h_3^z \sigma_3^z, \tag{81}$$

where the spins 1 and 3 are the focused spins and the spin 2 is the media spin (Fig. 2). We assume $J^x \geq J^y \geq J^z$. We solve the entanglement maximization problem about the spins 1 and 3 by fixing the temperature T and the field h^z_{media} on the spin 2. In order to solve this maximization problem numerically, we use the random search method and the Newton method together. According to Theorem 1, if the two spins were separated by two spins, there would always be a critical temperature above which the maximized entanglement is exactly equal to zero. In the present case, the focused two spins are separated by only one spin and therefore Theorem 1 does not apply; the elements $|F_1|$, $\sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}}$, $|F_2|$ and $\sqrt{p_{\uparrow\downarrow}p_{\downarrow\uparrow}}$ are shown to be of the same order in the same way as in the proof of Theorem 1. Therefore, it generally depends on the interaction Hamiltonian and the positions of the focused spins whether the critical temperature exists or not. As for the Hamiltonian (81), except the case of the Ising chain with $h^z_{\mathrm{media}} = 0$, we can prove [22] that the entanglement between the spins 1 and 3 can exist at any temperatures by letting $h_1^z = h_3^z \to \infty$.

In Fig. 3, we show the numerical results on the entanglement maximization. The main feature is that in a certain parameter region the maximizing local fields $h_{1\mathrm{op}}$ and $h_{3\mathrm{op}}$ are asymmetric to each other, namely $|h_{1\mathrm{op}}| \neq |h_{3\mathrm{op}}|$. In this region, the asymmetry must be essential to the enhancement of the entanglement. The asymmetry appears continuously (solid line in Fig. 3) or discontinuously (broken line in Fig. 3). Note that in these cases there is no critical temperature above which the maximized entanglement would be zero.

B. Analytical argument

Here, we argue the origin of the asymmetry for the XX model; the phase diagram of the other models are not much different from the XX model qualitatively. The main reason of the asymmetry is the strong dependence of the indirect interaction on the local fields. Then, we focus on the magnons which mediate the indirect interaction and discuss the effects of the local fields on the magnons. We show that the following three points affect the asymmetry:

1. The Boltzmann weights of the states with the magnons.
2. Localization of the magnons.
3. Suppression of the off-diagonal elements $\{F_1, F_2\}$ due to mixing of different magnon states.



(a) (b) (c) (d)

FIG. 3: The parameter ranges in which the asymmetry appears (a) in the case of $J^x = J^y = 1$ and (b) in the case of $J^x = 1, J^y = 0.5$. In the 'Asymmetric' phase, the optimizing fields $h_{1\mathrm{op}}$ and $h_{3\mathrm{op}}$ are asymmetric to each other as $|h_{1\mathrm{op}}| \neq |h_{3\mathrm{op}}|$, while they satisfy $|h_{1\mathrm{op}}| = |h_{3\mathrm{op}}|$ in the 'Symmetric' phase. On the solid line, the asymmetry appears continuously, while on the broken line the asymmetry appears discontinuously. The ratio $h_{1\mathrm{op}}/h_{3\mathrm{op}}$ is shown for (c) $h^z_{\mathrm{media}} = 4$ and (d) $T = 1.5$ on the chained lines of (a).

We define a magnon as a spin flip; a down spin in the background of up spins or an up spin in the background of down spins. For example, the magnon number is two for both of the states $|\uparrow\uparrow\uparrow\downarrow\downarrow\rangle$ and $|\uparrow\uparrow\downarrow\downarrow\downarrow\rangle$. In the XX three-spin chain, the number of the magnons can be either zero or one according to the above definition. The magnon does not exist in the states $|\uparrow_1\uparrow_2\uparrow_3\rangle$ and $|\downarrow_1\downarrow_2\downarrow_3\rangle$, nor the entanglement. This suggests that the magnons mediate the indirect interaction and is essential to the entanglement between the focused spins. Even if the magnons exist, however, the entanglement can be very small when the magnon is localized in one site. It is also possible that the off-diagonal elements $\{F_1, F_2\}$ in the density matrix (4) are suppressed by the mixture of the two kinds of magnon states. As shown in Eqs. (45) and (46), these elements are related to the correlation between the focused spins;

$$F_1 = \frac{\langle \sigma_2^x \sigma_{N-1}^x + \sigma_2^y \sigma_{N-1}^y \rangle_0}{4},$$
$$F_2 = \frac{\langle \sigma_2^x \sigma_{N-1}^x - \sigma_2^y \sigma_{N-1}^y \rangle_0}{4}. \tag{82}$$

Let us show an example; each of the two states

$$\frac{1}{\sqrt{2}}(|\uparrow_1\uparrow_2\downarrow_3\rangle + |\downarrow_1\uparrow_2\uparrow_3\rangle),\ \frac{1}{\sqrt{2}}(|\uparrow_1\uparrow_2\downarrow_3\rangle - |\downarrow_1\uparrow_2\uparrow_3\rangle), \tag{83}$$

has one magnon and its entanglement is maximum. However, the correlations which their magnons mediate are opposite to each other; the first state has $\langle \sigma_1^x \sigma_3^x \rangle = \langle \sigma_1^y \sigma_3^y \rangle = 1$, whereas the second one has $\langle \sigma_1^x \sigma_3^x \rangle = \langle \sigma_1^y \sigma_3^y \rangle = -1$. Then, the off-diagonal elements $\{F_1, F_2\}$ are also opposite to each other. Therefore, the entanglement is enhanced if we suppress the mixedness of the one state of (83).

Let us analytically show how the above three points affect the asymmetry. First, we consider the case of $T \simeq 0$ and $h^z_{\mathrm{media}} \gg J$, that is, the upper left corner of the phase diagram in Fig. 3. In this case, we have $h_{1\mathrm{op}} = h_{3\mathrm{op}} > 0$ in the 'Symmetric' phase and $h_{1\mathrm{op}}, h_{3\mathrm{op}} > 0$ but $h_{1\mathrm{op}} \neq h_{3\mathrm{op}}$ in the 'Asymmetric' phase. Let us consider the interaction J and the local fields $h_{1\mathrm{op}}, h_{3\mathrm{op}}$ on the focused spins as perturbation. Then, the four unperturbed ground states $|{\uparrow_1\uparrow_2\uparrow_3}\rangle$, $|{\uparrow_1\uparrow_2\downarrow_3}\rangle$, $|{\downarrow_1\uparrow_2\uparrow_3}\rangle$ and $|{\downarrow_1\uparrow_2\downarrow_3}\rangle$ are degenerate; we define these four states as $\{\psi^{(0)}_i\}$. Because we assume $h^z_{\mathrm{media}} \gg J$, we consider the mixing of only these four states. The first-order contribution of the excited states $|{\uparrow_1\downarrow_2\uparrow_3}\rangle$, $|{\uparrow_1\downarrow_2\downarrow_3}\rangle$, $|{\downarrow_1\downarrow_2\uparrow_3}\rangle$ and $|{\downarrow_1\downarrow_2\downarrow_3}\rangle$ to the above four unperturbed ground states are given in the forms

$$\begin{aligned}
&|\psi_1\rangle = |{\uparrow_1\uparrow_2\uparrow_3}\rangle,\ \epsilon_1 = -h_{1\mathrm{op}} - h_{3\mathrm{op}} - h^z_{\mathrm{media}},\\
&|\psi_2\rangle = \mathcal{K}_+\big(|{\uparrow_1\uparrow_2\downarrow_3}\rangle + \frac{\delta}{J}|{\uparrow_1\downarrow_2\uparrow_3}\rangle\big) + \mathcal{K}_-\big(|{\downarrow_1\uparrow_2\uparrow_3}\rangle + \frac{\delta}{J}|{\uparrow_1\downarrow_2\uparrow_3}\rangle\big),\\
&\quad \epsilon_2 = -\delta - \sqrt{(h_{1\mathrm{op}} - h_{3\mathrm{op}})^2 + \delta^2} - h^z_{\mathrm{media}},\\
&|\psi_3\rangle = \mathcal{K}_-\big(|{\uparrow_1\uparrow_2\downarrow_3}\rangle + \frac{\delta}{J}|{\uparrow_1\downarrow_2\uparrow_3}\rangle\big) - \mathcal{K}_+\big(|{\downarrow_1\uparrow_2\uparrow_3}\rangle + \frac{\delta}{J}|{\uparrow_1\downarrow_2\uparrow_3}\rangle\big),\\
&\quad \epsilon_3 = -\delta + \sqrt{(h_{1\mathrm{op}} - h_{3\mathrm{op}})^2 + \delta^2} - h^z_{\mathrm{media}},\\
&|\psi_4\rangle = |{\downarrow_1\uparrow_2\downarrow_3}\rangle + \frac{\delta}{J}|{\downarrow_1\downarrow_2\uparrow_3}\rangle + \frac{\delta}{J}|{\uparrow_1\downarrow_2\downarrow_3}\rangle,\ \epsilon_4 = h_{1\mathrm{op}} + h_{3\mathrm{op}} - h^z_{\mathrm{media}} + 2\delta,
\end{aligned} \tag{84}$$

where $\delta \equiv -J^2/(2h^z_{\mathrm{media}})$ and

$$\mathcal{K}_\pm \equiv \frac{1}{\sqrt{2}}\sqrt{1 \pm \frac{h_{1\mathrm{op}} - h_{3\mathrm{op}}}{\sqrt{(h_{1\mathrm{op}} - h_{3\mathrm{op}})^2 + \delta^2}}}. \tag{85}$$

We thereby calculate the elements F_1, $p_{\uparrow\uparrow}$ and $p_{\downarrow\downarrow}$ of the density matrix ρ_{12}, which contribute to the concurrence. First, the leading terms of $p_{\uparrow\uparrow}$ and $p_{\downarrow\downarrow}$ are obtained as follows:

$$p_{\uparrow\uparrow} = \frac{e^{-\beta\epsilon_1}}{Z_{\mathrm{tot}}} \text{ and } p_{\downarrow\downarrow} = \frac{e^{-\beta\epsilon_4}}{Z_{\mathrm{tot}}}. \tag{86}$$

Therefore, $\sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}}$ is equal to $e^{\beta(h^z_{\mathrm{media}}-\delta)}/Z_{\mathrm{tot}}$ and does not depend on $h_{1\mathrm{op}}$ and $h_{3\mathrm{op}}$. Then the entanglement depends on $h_{1\mathrm{op}}$ and $h_{3\mathrm{op}}$ mainly through F_1 as is shown in (7).

We obtain the leading term of F_1 as follows:

$$\begin{aligned}
F_1 = &\frac{e^{\beta(h^z_{\mathrm{media}}+\delta)}}{Z_{\mathrm{tot}}}\mathcal{K}_+\mathcal{K}_-\\
&\times\left(e^{\beta\sqrt{(h_{1\mathrm{op}}-h_{3\mathrm{op}})^2+\delta^2}} - e^{-\beta\sqrt{(h_{1\mathrm{op}}-h_{3\mathrm{op}})^2+\delta^2}}\right).
\end{aligned} \tag{87}$$

Let us see how F_1 depends on the asymmetry $h_{1\mathrm{op}} - h_{3\mathrm{op}}$. We consider the following three terms of F_1 separately:

$$\begin{aligned}
&\mathcal{K}_+\mathcal{K}_- = \frac{1}{2}\left(1 - \frac{(h_{1\mathrm{op}} - h_{3\mathrm{op}})^2}{(h_{1\mathrm{op}} - h_{3\mathrm{op}})^2 + \delta^2}\right),\\
&e^{\beta\sqrt{(h_{1\mathrm{op}}-h_{3\mathrm{op}})^2+\delta^2}} \text{ and } e^{-\beta\sqrt{(h_{1\mathrm{op}}-h_{3\mathrm{op}})^2+\delta^2}}.
\end{aligned} \tag{88}$$

We start from the point $h_{1\mathrm{op}} = h_{3\mathrm{op}}$ and gradually increase the asymmetry $h_{1\mathrm{op}} - h_{3\mathrm{op}}$ to see how the above three terms affect the entanglement through the element F_1. We argue that there are two effects competing with each other, which may yield the phase boundary in the upper left corner of the phase diagram in Fig. 3.

The two factors $\mathcal{K}_+$ and $\mathcal{K}_-$ on the right-hand side of the first term in (88) appear as coefficients in the states $|\psi_2\rangle$ and $|\psi_3\rangle$ in (84). As the asymmetry $h_{1\mathrm{op}} - h_{3\mathrm{op}}$ is increased in Eq. (84), the factor $\mathcal{K}_-$ decreases and therefore the states $|\psi_2\rangle$ and $|\psi_3\rangle$ become dominated by the states $|{\uparrow_1\uparrow_2\downarrow_3}\rangle$ and $|{\downarrow_1\uparrow_2\uparrow_3}\rangle$, respectively; the factor $\mathcal{K}_+$ decreases when the asymmetry is reversed to $h_{3\mathrm{op}} - h_{1\mathrm{op}}$. Then the magnon becomes localized on the site 1 or 3 as the asymmetry $h_{1\mathrm{op}} - h_{3\mathrm{op}}$ is increased. This is a negative effect on the entanglement; the entanglement would be suppressed if the magnon does not move around the system. Indeed, the element F_1 in (87) and hence the entanglement in (7) can decrease because of the decrease of the first term in (88).

The second and third terms in (88) are the Boltzmann weights of the states $|\psi_2\rangle$ and $|\psi_3\rangle$. As the asymmetry $h_{1\mathrm{op}} - h_{3\mathrm{op}}$ is increased, the second term increases while the third term decreases exponentially. The state $|\psi_2\rangle$ has positive transverse correlations $\langle\sigma_1^x\sigma_3^x\rangle$ and $\langle\sigma_1^y\sigma_3^y\rangle$, while the state $|\psi_3\rangle$ has negative ones. As has been shown in the example of (83), the increase and decrease of the respective Boltzmann weights enhance the transverse correlations. This is a positive effect on the entanglement; the transverse correlations can promote the off-diagonal elements $\{F_1, F_2\}$ in Eq. (82). Indeed, the element F_1 in (87) can increase because of the exponential increase and the decrease of the second and third terms in (88).

The above negative and positive effects of the asymmetry on F_1 and the entanglement compete with each other. In other words, in some cases the entanglement is maximized without the asymmetry, while in other cases it is maximized by introducing the asymmetry. This may be the reason for the phase boundary in the upper left corner of the phase diagram in Fig. 3.

Second, we consider the case $h_{\mathrm{media}} > T \gg 1$, that is, the phase of the asymmetry in the upper right area of the phase diagram in Fig. 3. We numerically obtained $h_{1\mathrm{op}} \gg J$, $h_{3\mathrm{op}} \ll -J$ and $|h_{1\mathrm{op}}| \neq |h_{3\mathrm{op}}|$ in this area. We show that the asymmetry indeed promotes the entanglement by starting from the point $h_{1\mathrm{op}} = -h_{3\mathrm{op}}$ and increasing the asymmetry $h_{1\mathrm{op}} + h_{3\mathrm{op}}$ gradually. We argue that the increase of the entanglement is due to delocalization of the magnon. In order to calculate the entanglement in the present case, we regard the interaction between the focused spins and the media spins as perturbation.

Because $|h_{1\mathrm{op}}| \gg 1$ and $|h_{3\mathrm{op}}| \gg 1$, we consider only the two states $|{\uparrow_1}{\uparrow_2}{\downarrow_3}\rangle$ and $|{\uparrow_1}{\downarrow_2}{\downarrow_3}\rangle$ as the unperturbed states. Then, these two states in the first-order perturbation are given by

$$\begin{aligned}|\psi_1\rangle &= |{\uparrow_1}{\uparrow_2}{\downarrow_3}\rangle + \frac{J}{-2(|h_{3\mathrm{op}}| + h^z_{\mathrm{media}})}|{\uparrow_1}{\downarrow_2}{\uparrow_3}\rangle + \frac{J^2}{4(|h_{3\mathrm{op}}| + h^z_{\mathrm{media}})(h_{1\mathrm{op}} + |h_{3\mathrm{op}}|)}|{\downarrow_1}{\uparrow_2}{\uparrow_3}\rangle,\\ \epsilon_1 &= -h_{1\mathrm{op}} - |h_{3\mathrm{op}}| - h^z_{\mathrm{media}}\end{aligned} \tag{89}$$

and

$$\begin{aligned}|\psi_2\rangle &= |{\uparrow_1}{\downarrow_2}{\downarrow_3}\rangle + \frac{J}{-2(h_{1\mathrm{op}} + h^z_{\mathrm{media}})}|{\downarrow_1}{\uparrow_2}{\downarrow_3}\rangle + \frac{J^2}{4(h_{1\mathrm{op}} + h^z_{\mathrm{media}})(h_{1\mathrm{op}} + |h_{3\mathrm{op}}|)}|{\downarrow_1}{\downarrow_2}{\uparrow_3}\rangle,\\ \epsilon_2 &= -h_{1\mathrm{op}} - |h_{3\mathrm{op}}| + h^z_{\mathrm{media}}.\end{aligned} \tag{90}$$

From these expressions, we can calculate $\sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}}$ as

$$\sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}} = e^{\beta(h_{1\mathrm{op}}+|h_{3\mathrm{op}}|)}\frac{J^2}{4(|h_{3\mathrm{op}}| + h^z_{\mathrm{media}})(h_{1\mathrm{op}} + h^z_{\mathrm{media}})}. \tag{91}$$

Similarly, the element F_1 is given by

$$F_1 = e^{\beta(h_{1\mathrm{op}}+|h_{3\mathrm{op}}|)}\left(\frac{J^2}{4(|h_{3\mathrm{op}}| + h^z_{\mathrm{media}})(h_{1\mathrm{op}} + |h_{3\mathrm{op}}|)}e^{\beta h^z_{\mathrm{media}}} + \frac{J^2}{4(h_{1\mathrm{op}} + h^z_{\mathrm{media}})(h_{1\mathrm{op}} + |h_{3\mathrm{op}}|)}e^{-\beta h^z_{\mathrm{media}}}\right). \tag{92}$$

To clarify the effect of the asymmetry $h_{1\mathrm{op}} + h_{3\mathrm{op}}$, we put

$$h_{1\mathrm{op}} = h_0 + \delta h \text{ and } h_{3\mathrm{op}} = -h_0 + \delta h, \tag{93}$$

where $h_0 \gg \delta h > 0$. Then, Eqs. (91) and (92) are recast into

$$\sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}} = e^{2\beta h_0}\frac{J^2}{4\big[(h^z_{\mathrm{media}} + h_0)^2 - \delta h^2\big]} \tag{94}$$

and

$$F_1 = e^{2\beta h_0}\left(\frac{J^2}{8h_0(h_0 + h^z_{\mathrm{media}} - \delta h)}e^{\beta h^z_{\mathrm{media}}} + \frac{J^2}{8h_0(h_0 + h^z_{\mathrm{media}} + \delta h)}e^{-\beta h^z_{\mathrm{media}}}\right). \tag{95}$$

The former $\sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}}$ does not depend on the symmetry δh up to the first order. On the other hand, the latter F_1 and hence the entanglement in (7) increase as δh is increased; the increase of the first term on the right-hand side of (95) excels the decrease of the second term because of the difference in the Boltzmann weights.

The increase of F_1 can be related to delocalization of the magnon. Since $|h_{1\mathrm{op}}| \gg 1$ and $|h_{N\mathrm{op}}| \gg 1$, the magnon is almost localized in the site 3 in $|\psi_1\rangle$ of (89) and in the site 1 in $|\psi_2\rangle$ of (90), but slightly delocalized owing to the perturbation terms. As we increase δh, the perturbation terms of $|\psi_1\rangle$ increase, but those of $|\psi_2\rangle$ decrease. Therefore, the magnon is delocalized more in $|\psi_1\rangle$, while it is localized more in $|\psi_2\rangle$. This corresponds to the increase of the first term and the decrease of the second term in (95). Because the Boltzmann weight of the state $|\psi_1\rangle$ is greater than that of $|\psi_2\rangle$, the magnon delocalization in $|\psi_1\rangle$ excels the magnon localization in $|\psi_2\rangle$. This corresponds to the increase of F_1.



FIG. 4: Schematic picture of four-spin chains. We define the spins 1 and 4 as the focused spins whereas the spins 2 and 3 as the media spins. The spins 1 and 4 indirectly interact with each other through the spins 2 and 3. In this system, the two focused spins are separated by two spins. Therefore, according to Theorem 1, there is a critical temperature above which the maximized entanglement is equal to zero.



FIG. 5: The phase diagrams for the four-spin chain, (a) in the case of $J^x = J^y = 1$ and (b) in the case of $J^x = 1, J^y = 0.5$. In the 'Asymmetric' phase, the maximizing fields $h_{1\rm op}$ and $h_{4\rm op}$ are asymmetric to each other as $|h_{1\rm op}| \neq |h_{4\rm op}|$, while they satisfy $|h_{1\rm op}| = |h_{4\rm op}|$ in the 'Symmetric' phase. The asymmetry appears continuously on the solid line, while it appears discontinuously on the broken line. The maximized entanglement vanishes beyond the chained line.

V. ENTANGLEMENT MAXIMIZATION IN FOUR-SPIN CHAINS

In the present section, we consider the maximization problem in four-spin chains (Fig. 4). As has been proved in Theorem 1, the four-spin chain is the shortest one in which the end-to-end entanglement cannot be generated in the high-temperature limit. We mainly discuss the critical temperature and its dependence on the interaction of the spins.

A. Numerical results

In the present section, we consider the XY spin chains given by the Hamiltonian

$$\begin{aligned} H_{\rm tot} = &\sum_{i=1}^{3} (J^x \sigma_i^x \sigma_{i+1}^x + J^y \sigma_i^y \sigma_{i+1}^y) \\ &+ h_1^z \sigma_1^z + h_4^z \sigma_4^z + h_{\rm media}^z (\sigma_2^z + \sigma_3^z). \end{aligned} \tag{96}$$

We solve the entanglement maximization problem about the focused spins 1 and 4 by fixing the temperature T and the field $h_{\rm media}^z$ on the media spins 2 and 3. In order to solve this maximization problem numerically, we used the random search method and the Newton method together. According to Theorem 1, there always exists a critical temperature above which the maximized entanglement is exactly equal to zero because the focused spins 1 and 4 are separated by two spins.

We show the phase diagram of the XX spin chain and the XY spin chain in Fig. 5. As in the case of the three spins, there is an 'Asymmetric' phase where the maximizing local field $|h_{1\text{op}}|$ is not equal to $|h_{4\text{op}}|$. Moreover, the maximized entanglement is equal to zero in a region. The qualitatively different behavior of the critical temperature between the XX and the XY chains is due to the conservation of the angular momentum in the z direction, as we will argue in Section V B.

B. Difference between the XX and the XY model

We discuss the behavior of the critical temperature in the XX and XY chains. In the XX chain, the critical temperature increases as the media field h^z_{media} is increased, while in the XY chain it does not. This difference is attributed to the conservation of the angular momentum in the z direction. In the XX chain, we can suppress the mixture of the states with more than two magnons by choosing the fields as

$$\begin{aligned} &h^z_1 = -h_0 \\ &h^z_i = h_0, \text{ for } i = 2, 3, \cdots N, \end{aligned} \tag{97}$$

and $h_0\beta \gg 1$. Then, the density matrix $e^{-\beta H_{\text{tot}}}$ is almost equivalent to the ground state of H_{tot} and the mixture of the other states can be suppressed exponentially by increasing h_0. The ground state is given by the following form;

$$\begin{aligned} &a_0|{\downarrow_1}{\uparrow_2}\cdots{\uparrow_N}\rangle + a_1|{\uparrow_1}{\downarrow_2}\cdots{\uparrow_N}\rangle + a_2|{\uparrow_1}{\uparrow_2}{\downarrow_3}\cdots{\uparrow_N}\rangle \\ &+\cdots + a_{N-1}|{\uparrow_1}{\uparrow_2}\cdots{\downarrow_N}\rangle, \end{aligned} \tag{98}$$

where we can calculate a_k by the kth order perturbation to have

$$a_k = O\left(\frac{J^k}{h_0^k}\right) \tag{99}$$

for $k = 1, 2, \cdots N-1$ with the factor h_0^k coming from the energy denominator. In the ground state (98), the element $p_{\downarrow\downarrow}$ is equal to zero because there is no state with more than one down spins in (98). The entanglement between the spins 1 and N exists because $|F_1| \propto |a_{N-1}| \propto |J/h_0|^{N-1} > 0$. The mixture of the excited states generally destroys the entanglement, but is suppressed exponentially because of the Boltzmann weights. The entanglement between the spins 1 and N thereby survives.

On the other hand, in the XY chains, we cannot control the number of the magnons in the ground state by increasing the media fields. Therefore, $p_{\uparrow\uparrow}p_{\downarrow\downarrow}$ is not zero in the ground state, which invalidates the argument for the XX model. This may account for the fact that the critical temperature does not increase as the media fields h^z_{media} are increased.

C. Calculation of the critical temperature

Let us calculate the critical temperature analytically. It is generally difficult to solve the entanglement maximization analytically. At the critical temperature, however, we numerically confirmed that the maximizing local fields take the form $h_{1\text{op}} = h_{4\text{op}} \to \infty$ in some region of the 'Symmetric' phases in Fig. 5. For the XX model, this happens everywhere in the 'Symmetric' phase in Fig. 5 (a), whereas for the XY model it happens only in the lower one of the two 'Symmetric' phases Fig. 5 (b). In such regions, we can derive the critical temperature analytically. In the 'Asymmetric' phases in Fig. 5, on the other hand, the maximizing local fields do not have simple forms.

After the calculation in Ref. [23], we obtain a necessary and sufficient condition (8) for the existence of the entanglement in the form of the following two inequalities:

$$\begin{aligned} &Z^2_{\text{tot}}(F_1^2 - p_{\uparrow\uparrow}p_{\downarrow\downarrow}) = \left[4\mathcal{S}_-\mathcal{S}_+\gamma\sinh\beta|\epsilon_3| + (1+\gamma^2)\sinh\beta|\epsilon_1|\right]^2 \\ &-Z_{\text{tot}}\left[(\gamma^4\mathcal{S}_-^2 + \mathcal{S}_+^2)e^{-\beta\epsilon_4} + (\mathcal{S}_-^2 + \gamma^4\mathcal{S}_+^2)e^{-\beta\epsilon_3} + 2\gamma^2\cosh\beta|\epsilon_1|\right] > 0, \end{aligned} \tag{100}$$

or

$$\begin{aligned} &Z^2_{\text{tot}}(F_2^2 - p_{\uparrow\downarrow}p_{\downarrow\uparrow}) = \left[2\mathcal{S}_-\mathcal{S}_+(1+\gamma^2)\sinh\beta|\epsilon_3| + 2\gamma\sinh\beta|\epsilon_1|\right] \\ &-\left[(\gamma^2\mathcal{S}_-^2 + \mathcal{S}_+^2)e^{-\beta\epsilon_4} + (\mathcal{S}_-^2 + \gamma^2\mathcal{S}_+^2)e^{-\beta\epsilon_3} + (\gamma^2+1)\cosh\beta|\epsilon_1|\right] > 0. \end{aligned} \tag{101}$$



(a) (b)

FIG. 6: Comparison of the phase boundary (a) in the case of $J^x = J^y = 1$ and (b) in the case of $J^x = 1, J^y = 0.5$. The critical temperature calculated according to the inequality (100) is indicated by the chained lines and the numerically calculated one by the solid lines, which are the same as in Fig. 5. The critical temperature calculated by the inequality (101) is also indicated by the broken line in (b). In the XX spin chain, the approximation gives almost the same values as the numerical ones in the 'symmetric' phase, while in the XY spin chain, the approximated values from the inequality (100) fit the numerical ones in the range of $0 \leq h^z_{\mathrm{media}} \leq 0.851$.

where J and γ are defined in (35),

$$\mathcal{S}_\pm \equiv \frac{1}{\sqrt{2}}\sqrt{1 \pm \frac{2h^z_{\mathrm{media}}}{\sqrt{4(h^z_{\mathrm{media}})^2 + (J^x - J^y)^2}}}, \tag{102}$$

and

$$\begin{aligned}\epsilon_1 &= -\epsilon_2 = -J^x - J^y,\\ \epsilon_3 &= -\epsilon_4 = -\sqrt{4(h^z_{\mathrm{media}})^2 + (J^x - J^y)^2},\\ Z_{\mathrm{tot}} &= e^{-\beta\epsilon_1} + e^{-\beta\epsilon_2} + e^{-\beta\epsilon_3} + e^{-\beta\epsilon_4}.\end{aligned} \tag{103}$$

In the case of the XX model with $\gamma = 0$, in particular, the above condition becomes simpler; because we have $F_2 = 0$ for $\gamma = 0$, the inequality (101) is never satisfied. The condition (100), on the other hand, reduces to

$$\sinh^2(\beta|\epsilon 1|) - Z_{\mathrm{tot}}e^{-\beta\epsilon_4} = \sinh^2(2\beta J) - Z_{\mathrm{tot}}e^{-2\beta h^z_{\mathrm{media}}} > 0, \tag{104}$$

or

$$\frac{e^{2\beta h^z_{\mathrm{media}}}\sinh^2(2\beta J)}{2\cosh(2\beta J) + 2\cosh(2\beta h^z_{\mathrm{media}})} > 1. \tag{105}$$

In Fig. 6, we compare the critical temperatures calculated according to the inequalities. (100) and (101) with the numerically calculated ones. The approximations generally give precise estimates for the XX model in the 'Symmetric' phase. For the XY model, the approximate estimate shown in Fig. 6 (b) is due to (100) in the range bounded by the broken line and is due to (101) in the range bounded by the thin solid line. The former gives good estimates in the range of $0 \leq h^z_{\mathrm{media}} \leq 0.851$.

VI. SUMMARY AND CONCLUSION

We have analytically and numerically studied the maximum value of the thermal entanglement between two spins which indirectly interact with each other. We showed the general theorem on its temperature dependence. We proved that the maximized entanglement is equal to zero above the critical temperature if the two spins are separated by two or more spins. This result tells us that as the temperature goes higher the thermal entanglement between distantly separated two spins cannot survive with any local controls.

Secondly, in the three-spin chains and the four-spin chains, we showed properties of the maximized entanglement which is calculated numerically. In the three-spin chains, we showed that the maximizing local fields are not symmetric as $h_{1\mathrm{op}} \neq h_{3\mathrm{op}}$ (Fig. 3) in some parameter regions. We have explained the asymmetry qualitatively and quantitatively. We attributed the appearance of the asymmetry to the effects on the magnons, which mediate the indirect interaction. In other words, the local fields affect the effective interaction between the focused spins, while the direct interaction is determined independently of the external fields. In the four-spin chains, we demonstrated that the maximized entanglement vanishes above the certain critical temperature. Because of the difference in the symmetric properties between the XX and XY spin chains, the dependence of the critical temperatures on the media fields are qualitatively different between the two systems. We calculated the critical temperature analytically in some parameter regions.

In conclusion, we have clarified several properties of the entanglement which is generated from the indirect interaction. Our study has given one of the general limits for the entanglement generation. We have also shown some properties of the external fields which is closely related to the entanglement enhancement. However, there are many problems to be solved on the general relationship between the external fields and the entanglement enhancement. In particular, we have not shown the properties of the multipartite entanglement. It is obvious that the external fields also have effect on qualitative behaviors of the multipartite entanglement; the most famous one is the quantum phase transition of the transverse Ising model [24]. In future, we plan to investigate the general properties of the enhancement of the multipartite entanglement by external fields.

ACKNOWLEDGMENT

The present author is grateful to Professor Naomichi Hatano for helpful discussions and comments.

Appendix A: In the case $h_{1\mathrm{op}} - h_{N\mathrm{op}} = O(\beta^{-\tilde{\kappa}})$ with $\tilde{\kappa} \le 0$ in the case (b)

Here, we discuss the case of $h_{1\mathrm{op}} - h_{N\mathrm{op}} = O(\beta^{-\tilde{\kappa}})$ with $\tilde{\kappa} \le 0$ in the case (b). In this case, we cannot consider the unperturbed states $|\uparrow_1\downarrow_N\rangle \otimes |\psi^n_{\mathrm{media}}\rangle$ and $|\downarrow_1\uparrow_N\rangle \otimes |\psi^n_{\mathrm{media}}\rangle$ independently because their eigenvalues are almost degenerate. Then, the magnitudes of the elements F_1 and F_2 can be different from the ones in (15). We can still apply the same calculation to the other parameters $\{P^{\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}, P^{\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}\}$ as in the case $\tilde{\kappa} > 0$; they are of order $\beta^{2\kappa_1+2\kappa_N}$.

We here prove that F_1^2 and F_2^2 are of order $\beta^{2\kappa_1+2\kappa_N}$ or higher. In order to prove this, we separate $H_1 + H_N$ as follows;

$$
\begin{aligned}
H_1 + H_N &= H_{\mathrm{LO}} + \delta H_{\mathrm{LO}}, \\
H_{\mathrm{LO}} &= \frac{1}{2}(h_{1\mathrm{op}} + h_{N\mathrm{op}} + h_0)\sigma_1^z + \frac{1}{2}(h_{1\mathrm{op}} + h_{N\mathrm{op}} - h_0)\sigma_N^z \\
&= -(h_{1\mathrm{op}} + h_{N\mathrm{op}})\big(|\uparrow_1\uparrow_N\rangle\langle\uparrow_1\uparrow_N| - |\downarrow_1\downarrow_N\rangle\langle\downarrow_1\downarrow_N|\big) - h_0\big(|\uparrow_1\downarrow_N\rangle\langle\uparrow_1\downarrow_N| - |\downarrow_1\uparrow_N\rangle\langle\downarrow_1\uparrow_N|\big), \\
\delta H_{\mathrm{LO}} &= -(h_{1\mathrm{op}} - h_{N\mathrm{op}} - h_0)\big(|\uparrow_1\downarrow_N\rangle\langle\uparrow_1\downarrow_N| - |\downarrow_1\uparrow_N\rangle\langle\downarrow_1\uparrow_N|\big), \\
&= \frac{1}{2}(h_{1\mathrm{op}} - h_{N\mathrm{op}} - h_0)(\sigma_1^z - \sigma_N^z), \qquad \text{(A1)}
\end{aligned}
$$

where we define as $h_0 = O(\beta^{\tilde{\kappa_0}})$ with $0 < \tilde{\kappa_0} < 1$, and regard $\beta\delta H_{\mathrm{LO}}$ as perturbation. The unperturbed density matrix $\tilde{\rho}^{(0)}_{\mathrm{tot}}$ is given by

$$
\tilde{\rho}^{(0)}_{\mathrm{tot}} = e^{-\beta\tilde{H}^{(0)}_{\mathrm{tot}}}, \qquad \text{(A2)}
$$

where $\tilde{H}^{(0)}_{\mathrm{tot}}$ is defined by

$$
\tilde{H}^{(0)}_{\mathrm{tot}} = H_{\mathrm{couple}} + H_{\mathrm{media}} + H_{\mathrm{LO}}, \qquad \text{(A3)}
$$

where H_{media} and H_{couple} are given in (19). Because $h_0 = O(\beta^{-\tilde{\kappa}_0})$ with $0 < \tilde{\kappa}_0 < 1$, the magnitudes of the unperturbed elements $\{F_1^{(0)}, F_2^{(0)}\}$ of $\tilde{\rho}^{(0)}_{\mathrm{tot}}$ are given by

$$
O(\beta^{\kappa_1+\kappa_N+\kappa_0'}) \text{ and } O(\beta^{\kappa_1+\kappa_N+\kappa_0}), \qquad \text{(A4)}
$$

where

$$
\kappa_0 = \min(\kappa_N, 1) \text{ and } \kappa_0' = \min(\kappa_N, \tilde{\kappa}_0, 1). \qquad \text{(A5)}
$$

The density matrix in the first-order perturbation is given by

$$\rho_{\rm tot}=\frac{1}{Z^{(0)}+\delta Z}\bigg(e^{-\beta\tilde{H}^{(0)}_{\rm tot}}+\beta\int_0^1 e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}\delta H_{\rm LO}e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}dx\bigg), \tag{A6}$$

where $Z^{(0)}$ is the partition function of $e^{-\beta\tilde{H}^{(0)}_{\rm tot}}$, while $Z^{(0)}+\delta Z$ is the partition function of $e^{-\beta H_{\rm tot}}$. The elements F_1 and F_2 are given by

$$\begin{aligned}F_1&={\rm tr}_{1N}\langle\uparrow_1\downarrow_N|\rho_{\rm tot}|\downarrow_1\uparrow_N\rangle,\\ F_2&={\rm tr}_{1N}\langle\uparrow_1\uparrow_N|\rho_{\rm tot}|\downarrow_1\downarrow_N\rangle,\end{aligned} \tag{A7}$$

where ${\rm tr}_{1N}$ denotes the trace operation on the spins except the focused spins 1 and N. The first-order perturbations of the elements F_1 and F_2 are

$$\begin{aligned}-\beta(h_{1\rm op}-h_{N\rm op}-h_0){\rm tr}_{1N}\int_0^1 dx\bigg(&\langle\uparrow_1\downarrow_N|e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}|\uparrow_1\downarrow_N\rangle\langle\uparrow_1\downarrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle\\ &-\langle\uparrow_1\downarrow_N|e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle\langle\downarrow_1\uparrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle\bigg)\end{aligned} \tag{A8}$$

and

$$\begin{aligned}-\beta(h_{1\rm op}-h_{N\rm op}-h_0){\rm tr}_{1N}\int_0^1 dx\bigg(&\langle\uparrow_1\uparrow_N|e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}|\uparrow_1\downarrow_N\rangle\langle\uparrow_1\downarrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\downarrow_N\rangle\\ &-\langle\uparrow_1\uparrow_N|e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle\langle\downarrow_1\uparrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\downarrow_N\rangle\bigg),\end{aligned} \tag{A9}$$

respectively.

In order to estimate the order of the first-order perturbations of F_1 and F_2, we introduce $\{|\psi^{(0),n,\xi}_{{\rm tot},\xi}\rangle\}$ and $\{E^{(0),n,\xi}_{\rm tot}\}$ as defined in Eqs. (23) and (24), but for $\tilde{H}^{(0)}_{\rm tot}$. First, we have

$$\langle\uparrow_1\downarrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle=\sum_{\substack{n=1\\ \xi=\uparrow\uparrow,\uparrow\downarrow,\downarrow\uparrow,\downarrow\downarrow}}^{2^{N-2}} e^{-\beta(1-x)E^{(0),n,\xi}_{\rm tot}}|\psi^{(0),n,\xi}_{{\rm media},\uparrow\downarrow}\rangle\langle\psi^{(0),n,\xi}_{{\rm media},\downarrow\uparrow}|. \tag{A10}$$

From the calculation in Ref. [20], we obtain

$$\Big|\Big||\psi^{(0),n,\xi}_{{\rm media},\uparrow\downarrow}\rangle\langle\psi^{(0),n,\xi}_{{\rm media},\downarrow\uparrow}|\Big|\Big|=O(\beta^{\kappa_1+\kappa_N}). \tag{A11}$$

Therefore, we have

$$\frac{1}{Z^{(0)}(x)}\big|\big|\langle\uparrow_1\downarrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle\big|\big|=O(\beta^{\kappa_1+\kappa_N}), \tag{A12}$$

where

$$Z^{(0)}(x)\equiv{\rm tr}\big(e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}\big). \tag{A13}$$

Similarly, we obtain

$$\begin{aligned}\frac{1}{Z^{(0)}(x)}\big|\big|\langle\uparrow_1\uparrow_N|e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}|\uparrow_1\downarrow_N\rangle\big|\big|&=O(\beta^{\kappa_N}),\\ \frac{1}{Z^{(0)}(1-x)}\big|\big|\langle\uparrow_1\downarrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\downarrow_N\rangle\big|\big|&=O(\beta^{\kappa_1}),\\ \frac{1}{Z^{(0)}(x)}\big|\big|\langle\uparrow_1\uparrow_N|e^{-\beta x\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle\big|\big|&=O(\beta^{\kappa_1}),\\ \frac{1}{Z^{(0)}(1-x)}\big|\big|\langle\downarrow_1\uparrow_N|e^{-\beta(1-x)\tilde{H}^{(0)}_{\rm tot}}|\downarrow_1\uparrow_N\rangle\big|\big|&=O(\beta^{\kappa_N}).\end{aligned} \tag{A14}$$

As a result, we obtain the first-order perturbations of F_1 and F_2 as

$$O(\beta^{\kappa_1+\kappa_N+1-\tilde{\kappa}_0}), \tag{A15}$$

where we utilized $\beta(h_{1\mathrm{op}}-h_{N\mathrm{op}}-h_0)=O(\beta^{1-\tilde{\kappa}_0})$ and

$$\frac{Z^{(0)}(x)Z^{(0)}(1-x)}{Z^{(0)}+\delta Z}=O(\beta^0). \tag{A16}$$

We can similarly calculate higher-order perturbations of F_1 and F_2 to see that they are of order higher than (A15). Thus, in the case $h_{1\mathrm{op}}-h_{N\mathrm{op}}=O(\beta^{-\tilde{\kappa}})$ with $\tilde{\kappa}\leq 0$, F_1^2 and F_2^2 are of order higher than $\{P^{\uparrow\uparrow}_{\uparrow\downarrow,\downarrow\uparrow}, P^{\uparrow\uparrow}_{\uparrow\uparrow,\downarrow\downarrow}\}$.

---

[1] E. Schrödinger, Naturwiss. **23**, 807 (1935).
[2] R. Horodecki, P. Horodecki, M. Horodecki and K. Horodecki, Rev. Mod. Phys. **81**, 865 (2009).
[3] M. A. Nielsen and I. L. Chuang, *Quantum computation and Quantum information* (Cambridge University Press, Cambridge, 2000).
[4] C. H. Bennett and D. P. DiVincenzo, Nature (London) **404**, 247 (2000).
[5] M. C. Arnesen, S. Bose and V. Vedral, Phys. Rev. Lett. **87**, 017901 (2001).
[6] L. Amico, R. Fazio, A. Osterloh and V. Vedral, Rev. Mod. Phys. **80**, 517 (2008).
[7] D. Gunlycke, V. M. Kendon, V. Vedral and S. Bose, Phys. Rev. A **64**, 042302 (2001).
[8] T. J. Osborne and M. A. Nielsen, Phys. Rev. A **66**, 032110 (2002).
[9] D. Markham, J. Anders, V. Vedral and M. Murao, Europhys. Lett. **81**, 40006 (2008)
[10] Y. Nakata, D. Markham and M. Murao, Phys. Rev. A **79**, 042313 (2009).
[11] G. L. Kamta and A. F. Starace, Phys. Rev. Lett. **88**, 107901 (2002).
[12] X. Wang, Phys. Lett. A **281**, 101 (2001).
[13] M. Fujinaga and N. Hatano, J. Phys. Soc. Jpn. **76**, 094001 (2007).
[14] T. Kuwahara and N. Hatano, Phys. Rev. A **83**, 062311 (2011)
[15] L. Campos Venuti, C. Degli Esposti Boschi and M. Roncaglia, Phys. Rev. Lett. **96**, 247206 (2006).
[16] A. Ferreira, and J. M. Lopes dos Santos, Phys. Rev. A **77**, 034301 (2008).
[17] T. J. G. Apollaro, A. Cuccoli, A. Fubini, F. Plastina, and P. Verrucchi, Phys. Rev. A **77**, 062314 (2008)
[18] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
[19] K. Życzkowski, P. Horodecki, A. Sanpera and M. Lewenstein, Phys. Rev. A **58**, 883 (1998).
[20] Section I of the supplementary material.
[21] Section II of the supplementary material.
[22] Section III of the supplementary material.
[23] Section IV of the supplementary material.
[24] T. R. De Oliveira, G. Rigolin, M.C. De Oliveira and E. Miranda, Phy. Rev. Lett. **97**, 170401 (2006)

Supplementary material for the article “General theory of entanglement enhancement by external fields in spin chains”

In the present supplementary material, we give detailed calculations which cannot be provided in the main text of the article “General theory of entanglement enhancement by external fields in spin chains”.

S.1: Calculation in the case (b)

Here, we derive the approximated form of $|\psi_{\mathrm{tot}}^{n_0,\uparrow\uparrow}\rangle$ in Eq. (23) in the main text, to obtain Eqs. (36)–(39). In order to calculate perturbation of $|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle$, we employ the general perturbation theory,

$$|\psi\rangle = |\psi^{(0)}\rangle + \big(I_{\mathrm{tot}} - |\psi^{(0)}\rangle\langle\psi^{(0)}|\big)\frac{1}{E^{(0)} - H^{(0)}}\delta H|\psi\rangle, \tag{S.1}$$

where $H^{(0)}$ is the unperturbed Hamiltonian, $|\psi^{(0)}\rangle$ and $E^{(0)}$ are the unperturbed eigenstate and the unperturbed eigenvalue, respectively, and δH is the perturbative Hamiltonian. We calculate each element of $|\psi_{\mathrm{tot}}^{n_0,\uparrow\uparrow}\rangle$, namely $\{|\psi_{\mathrm{media},\xi}^{n_0,\uparrow\uparrow}\rangle\}$ in Eq. (23) with $\xi = \uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$. We first calculate $|\psi_{\mathrm{media},\downarrow\uparrow}^{n_0,\uparrow\uparrow}\rangle$, which is given by the first-order perturbation;

$$\begin{aligned}
&|\psi_{\mathrm{media},\downarrow\uparrow}^{n_0,\uparrow\uparrow}\rangle = |{\downarrow_1}{\uparrow_N}\rangle\langle{\downarrow_1}{\uparrow_N}| \otimes I_{\mathrm{media}} \\
&\quad\times \Big[I_{\mathrm{tot}} - \big(|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle\big)\big(\langle{\uparrow_1}{\uparrow_N}| \otimes \langle\psi_{\mathrm{media}}^{n_0}|\big)\Big]\frac{1}{E^{(0)} - H^{(0)}}H_{\mathrm{couple}}|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle \\
&= \sum_{n=1}^{2^{N-2}} |{\downarrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n}\rangle \frac{\big(\langle{\downarrow_1}{\uparrow_N}| \otimes \langle\psi_{\mathrm{media}}^{n}|\big)H_{\mathrm{couple}}\big(|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle\big)}{E_{\mathrm{media}}^{n_0} - h_{1\mathrm{op}} - h_{N\mathrm{op}} - (E_{\mathrm{media}}^{n} + h_{1\mathrm{op}} - h_{N\mathrm{op}})},
\end{aligned} \tag{S.2}$$

where we put $E^{(0)}$ to $E_{\mathrm{media}}^{n_0} - h_{1\mathrm{op}} - h_{N\mathrm{op}}$, $H^{(0)}$ to $H_1 + H_N + H_{\mathrm{media}}$, $|\psi^{(0)}\rangle$ to $|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle$ and δH to H_{couple} in Eq. (S.1). As has been stated, we assume $h_{1\mathrm{op}} = O(\beta^{-\kappa_1})$ and $h_{N\mathrm{op}} = O(\beta^{-\kappa_N})$ with $\kappa_1, \kappa_N > 0$. On the other hand, the eigenvalues $\{E_{\mathrm{media}}^{n}\}$ are of order β^0 because the media-spin Hamiltonian H_{media} is fixed. Therefore, Eq. (S.2) can be approximated by

$$\begin{aligned}
&|\psi_{\mathrm{media},\downarrow\uparrow}^{n_0,\uparrow\uparrow}\rangle \\
&= \sum_{n=1}^{2^{N-2}} |{\downarrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n}\rangle \frac{\big(\langle{\downarrow_1}{\uparrow_N}| \otimes \langle\psi_{\mathrm{media}}^{n}|\big)H_{\mathrm{couple}}\big(|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle\big)}{-2h_{1\mathrm{op}}}\Big(1 + O(\beta^{\kappa_1})\Big).
\end{aligned} \tag{S.3}$$

We can sum the leading term over the label n to obtain

$$|\psi_{\mathrm{media},\downarrow\uparrow}^{n_0,\uparrow\uparrow}\rangle = \frac{1}{-2h_{1\mathrm{op}}}\big(|{\downarrow_1}{\uparrow_N}\rangle\langle{\downarrow_1}{\uparrow_N}| \otimes I_{\mathrm{media}}\big)\Big(H_{\mathrm{couple}}|{\uparrow_1}{\uparrow_N}\rangle \otimes |\psi_{\mathrm{media}}^{n_0}\rangle + O(\beta^{\kappa_1})\Big), \tag{S.4}$$

where

$$\sum_{n=1}^{2^{N-2}} |\psi_{\mathrm{media}}^{n}\rangle\langle\psi_{\mathrm{media}}^{n}| = I_{\mathrm{media}} \tag{S.5}$$

is the identity operator in the whole space of the media spins. Using Eqs. (21) and (34) in Eq. (S.3), we have

$$\begin{aligned}
|\psi_{\mathrm{media},\downarrow\uparrow}^{n_0,\uparrow\uparrow}\rangle = \frac{J}{-2h_{1\mathrm{op}}}|{\downarrow_1}{\uparrow_N}\rangle \otimes \Big(&\gamma s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}_{n_0}^{\uparrow\uparrow}\rangle|{\uparrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}_{n_0}^{\uparrow\downarrow}\rangle|{\downarrow_{N-1}}\rangle \\
&+ u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}_{n_0}^{\downarrow\uparrow}\rangle|{\uparrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}_{n_0}^{\downarrow\downarrow}\rangle|{\downarrow_{N-1}}\rangle + O(\beta^{\kappa_1})\Big).
\end{aligned} \tag{S.6}$$

Second, the leading term of $|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\uparrow\downarrow}\rangle$ is similarly given as follows:

$$
\begin{aligned}
&|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\uparrow\downarrow}\rangle \\
=&\sum_{n=1}^{2^{N-2}} |{\uparrow_1\downarrow_N}\rangle \otimes |\psi^{n}_{\mathrm{media}}\rangle \frac{\big(\langle{\uparrow_1\downarrow_N}| \otimes \langle\psi^{n}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big)}{-2h_{N\mathrm{op}}} \Big(1+O(\beta^{\kappa_N})\Big) \\
=&\frac{1}{-2h_{N\mathrm{op}}}\big(|{\uparrow_1\downarrow_N}\rangle\langle{\uparrow_1\downarrow_N}| \otimes I_{\mathrm{media}}\big)\Big(H_{\mathrm{couple}}|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle + O(\beta^{\kappa_N})\Big) \\
=&\frac{J}{-2h_{N\mathrm{op}}}|{\uparrow_1\downarrow_N}\rangle \otimes \Big(\gamma s_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + t_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle \\
&+\gamma u_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + w_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + O(\beta^{\kappa_N})\Big).
\end{aligned}
\tag{S.7}
$$

Third, we calculate $|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\downarrow\downarrow}\rangle$, which is given by the second-order perturbation;

$$
\begin{aligned}
&|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\downarrow\downarrow}\rangle = \sum_{n,n'=1}^{2^{N-2}} |{\downarrow_1\downarrow_N}\rangle \otimes |\psi^{n}_{\mathrm{media}}\rangle \\
&\Bigg[\frac{\big(\langle{\downarrow_1\downarrow_N}| \otimes \langle\psi^{n}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\uparrow_1\downarrow_N}\rangle \otimes |\psi^{n'}_{\mathrm{media}}\rangle\big)}{E^{n_0}_{\mathrm{media}} - h_{1\mathrm{op}} - h_{N\mathrm{op}} - (E^{n}_{\mathrm{media}} + h_{1\mathrm{op}} + h_{N\mathrm{op}})} \cdot \frac{\big(\langle{\uparrow_1\downarrow_N}| \otimes \langle\psi^{n'}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big)}{E^{n_0}_{\mathrm{media}} - h_{1\mathrm{op}} - h_{N\mathrm{op}} - (E^{n'}_{\mathrm{media}} - h_{1\mathrm{op}} + h_{N\mathrm{op}})} \\
&+\frac{\big(\langle{\downarrow_1\downarrow_N}| \otimes \langle\psi^{n}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\downarrow_1\uparrow_N}\rangle \otimes |\psi^{n'}_{\mathrm{media}}\rangle\big)}{E^{n_0}_{\mathrm{media}} - h_{1\mathrm{op}} - h_{N\mathrm{op}} - (E^{n}_{\mathrm{media}} + h_{1\mathrm{op}} + h_{N\mathrm{op}})} \cdot \frac{\big(\langle{\downarrow_1\uparrow_N}| \otimes \langle\psi^{n'}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big)}{E^{n_0}_{\mathrm{media}} - h_{1\mathrm{op}} - h_{N\mathrm{op}} - (E^{n'}_{\mathrm{media}} + h_{1\mathrm{op}} - h_{N\mathrm{op}})}\Bigg].
\end{aligned}
\tag{S.8}
$$

By utilizing the assumptions $h_{1\mathrm{op}} = O(\beta^{-\kappa_1}) > 0$, $h_{N\mathrm{op}} = O(\beta^{-\kappa_N}) > 0$ with $\kappa_1 \geq \kappa_N > 0$ and $E^{n}_{\mathrm{media}} = O(\beta^{0})$, we can approximate (S.8) as

$$
\begin{aligned}
|\psi^{n_0,\uparrow\uparrow}_{\mathrm{media},\downarrow\downarrow}\rangle = \sum_{n,n'=1}^{2^{N-2}} |{\downarrow_1\downarrow_N}\rangle \otimes |\psi^{n}_{\mathrm{media}}\rangle &\Bigg[\frac{\big(\langle{\downarrow_1\downarrow_N}| \otimes \langle\psi^{n}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\uparrow_1\downarrow_N}\rangle \otimes |\psi^{n'}_{\mathrm{media}}\rangle\big)}{-2(h_{1\mathrm{op}} + h_{N\mathrm{op}})} \\
&\times \frac{\big(\langle{\uparrow_1\downarrow_N}| \otimes \langle\psi^{n'}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big)}{-2h_{N\mathrm{op}}}\Big(1+O(\beta^{\kappa_N})\Big) \\
&+ \frac{\big(\langle{\downarrow_1\downarrow_N}| \otimes \langle\psi^{n}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\downarrow_1\uparrow_N}\rangle \otimes |\psi^{n'}_{\mathrm{media}}\rangle\big)}{-2(h_{1\mathrm{op}} + h_{N\mathrm{op}})} \\
&\times \frac{\big(\langle{\downarrow_1\uparrow_N}| \otimes \langle\psi^{n'}_{\mathrm{media}}|\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big)}{-2h_{1\mathrm{op}}}\Big(1+O(\beta^{\kappa_1})\Big)\Bigg].
\end{aligned}
\tag{S.9}
$$

We can calculate the leading terms of (S.9) as

$$
\begin{aligned}
&\frac{\big(|{\downarrow_1\downarrow_N}\rangle\langle{\downarrow_1\downarrow_N}| \otimes I_{\mathrm{media}}\big) H_{\mathrm{couple}}\big(|{\uparrow_1\downarrow_N}\rangle\langle{\uparrow_1\downarrow_N}| \otimes I_{\mathrm{media}}\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big)}{4h_{N\mathrm{op}}(h_{1\mathrm{op}} + h_{N\mathrm{op}})} \\
&+\frac{\big(|{\downarrow_1\downarrow_N}\rangle\langle{\downarrow_1\downarrow_N}| \otimes I_{\mathrm{media}}\big) H_{\mathrm{couple}}\big(|{\downarrow_1\uparrow_N}\rangle\langle{\downarrow_1\uparrow_N}| \otimes I_{\mathrm{media}}\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big)}{4h_{1\mathrm{op}}(h_{1\mathrm{op}} + h_{N\mathrm{op}})}.
\end{aligned}
\tag{S.10}
$$

Upon using Eqs. (21) and (34), the numerator of the first term in (S.10) reduces to

$$
\begin{aligned}
&\big(|{\downarrow_1\downarrow_N}\rangle\langle{\downarrow_1\downarrow_N}| \otimes I_{\mathrm{media}}\big) H_{\mathrm{couple}}\big(|{\uparrow_1\downarrow_N}\rangle\langle{\uparrow_1\downarrow_N}| \otimes I_{\mathrm{media}}\big) H_{\mathrm{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\mathrm{media}}\rangle\big) \\
=&\big(|{\downarrow_1\downarrow_N}\rangle\langle{\downarrow_1\downarrow_N}| \otimes I_{\mathrm{media}}\big) H_{\mathrm{couple}} J \\
\times&\Big[|{\uparrow_1\downarrow_N}\rangle \otimes \big(\gamma s_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + t_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + \gamma u_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + w_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle\big)\Big] \\
=&J^2|{\downarrow_1\downarrow_N}\rangle \otimes \big(\gamma^2 s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + \gamma u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle\big).
\end{aligned}
\tag{S.11}
$$

Similarly, the numerator of the second term in (S.10) reduces to

$$\begin{aligned}
&\big(|{\downarrow_1\downarrow_N}\rangle\langle{\downarrow_1\downarrow_N}| \otimes I_{\text{media}}\big) H_{\text{couple}} \big(|{\downarrow_1\uparrow_N}\rangle\langle{\downarrow_1\uparrow_N}| \otimes I_{\text{media}}\big) H_{\text{couple}} \big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n_0}_{\text{media}}\rangle\big) \\
=&\big(|{\downarrow_1\downarrow_N}\rangle\langle{\downarrow_1\downarrow_N}| \otimes I_{\text{media}}\big) H_{\text{couple}} J \\
&\times \Big[|{\downarrow_1\uparrow_N}\rangle \otimes \big(\gamma s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle\big)\Big] \\
=&J^2|{\downarrow_1\downarrow_N}\rangle \otimes \big(\gamma^2 s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + \gamma u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle\big),
\end{aligned} \tag{S.12}$$

which is equal to (S.11). As a result, we arrive at

$$\begin{aligned}
|\psi^{n_0,\uparrow\uparrow}_{\text{media},\downarrow\downarrow}\rangle =&\frac{J^2}{4h_{1\text{op}}h_{N\text{op}}}\Big(\gamma^2 s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle \\
&+ \gamma u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + O(\beta^{\kappa_N})\Big).
\end{aligned} \tag{S.13}$$

Next, we calculate each element of $|\psi^{n_0,\uparrow\downarrow}_{\text{tot}}\rangle$, namely $\{|\psi^{n_0,\uparrow\downarrow}_{\text{tot},\xi}\rangle\}$ in Eq. (23) with $\xi =\uparrow\uparrow, \uparrow\downarrow, \downarrow\uparrow, \downarrow\downarrow$. By similar calculations, we obtain

$$\begin{aligned}
|\psi^{n_0,\uparrow\downarrow}_{\text{media},\uparrow\uparrow}\rangle =&\frac{J}{2h_{N\text{op}}}\Big(s_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma t_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle \\
&+ u_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma w_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + O(\beta^{\kappa_N})\Big) \\
|\psi^{n_0,\uparrow\downarrow}_{\text{media},\downarrow\downarrow}\rangle =&\frac{J}{-2h_{1\text{op}}}\Big(\gamma s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + \gamma t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle \\
&+ u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + O(\beta^{\kappa_1})\Big).
\end{aligned} \tag{S.14}$$

The element $|\psi^{n_0,\uparrow\downarrow}_{\text{media},\downarrow\uparrow}\rangle$ can be calculated from

$$\begin{aligned}
&|\psi^{n_0,\uparrow\downarrow}_{\text{media},\downarrow\uparrow}\rangle = \sum_{n,n'=1}^{2^{N-2}} |{\downarrow_1\downarrow_N}\rangle \otimes |\psi^{n}_{\text{media}}\rangle \\
&\Bigg[\frac{\big(\langle{\downarrow_1\uparrow_N}| \otimes \langle\psi^{n}_{\text{media}}|\big) H_{\text{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n'}_{\text{media}}\rangle\big)}{E^{n_0}_{\text{media}} - h_{1\text{op}} + h_{N\text{op}} - (E^{n}_{\text{media}} + h_{1\text{op}} - h_{N\text{op}})} \cdot \frac{\big(\langle{\uparrow_1\uparrow_N}| \otimes \langle\psi^{n'}_{\text{media}}|\big) H_{\text{couple}}\big(|{\uparrow_1\downarrow_N}\rangle \otimes |\psi^{n_0}_{\text{media}}\rangle\big)}{E^{n_0}_{\text{media}} - h_{1\text{op}} + h_{N\text{op}} - (E^{n'}_{\text{media}} - h_{1\text{op}} - h_{N\text{op}})} \\
&+ \frac{\big(\langle{\downarrow_1\uparrow_N}| \otimes \langle\psi^{n}_{\text{media}}|\big) H_{\text{couple}}\big(|{\downarrow_1\downarrow_N}\rangle \otimes |\psi^{n'}_{\text{media}}\rangle\big)}{E^{n_0}_{\text{media}} - h_{1\text{op}} + h_{N\text{op}} - (E^{n}_{\text{media}} + h_{1\text{op}} - h_{N\text{op}})} \cdot \frac{\big(\langle{\downarrow_1\downarrow_N}| \otimes \langle\psi^{n'}_{\text{media}}|\big) H_{\text{couple}}\big(|{\uparrow_1\downarrow_N}\rangle \otimes |\psi^{n_0}_{\text{media}}\rangle\big)}{E^{n_0}_{\text{media}} - h_{1\text{op}} + h_{N\text{op}} - (E^{n'}_{\text{media}} + h_{1\text{op}} + h_{N\text{op}})}\Bigg].
\end{aligned} \tag{S.15}$$

By utilizing the assumptions $h_{1\text{op}} = O(\beta^{-\kappa_1})$, $h_{N\text{op}} = O(\beta^{-\kappa_N})$ with $\kappa_1 \geq \kappa_N > 0$, $h_{1\text{op}} - h_{N\text{op}} = O(\beta^{-\tilde{\kappa}})$ with $\tilde{\kappa} > 0$ and $E^{n}_{\text{media}} = O(\beta^0)$, we can approximate (S.15) as

$$\begin{aligned}
|\psi^{n_0,\uparrow\downarrow}_{\text{media},\downarrow\uparrow}\rangle = \sum_{n,n'=1}^{2^{N-2}} |{\downarrow_1\downarrow_N}\rangle \otimes |\psi^{n}_{\text{media}}\rangle &\Bigg[\frac{\big(\langle{\downarrow_1\uparrow_N}| \otimes \langle\psi^{n}_{\text{media}}|\big) H_{\text{couple}}\big(|{\uparrow_1\uparrow_N}\rangle \otimes |\psi^{n'}_{\text{media}}\rangle\big)}{-2(h_{1\text{op}} - h_{N\text{op}})} \\
&\times \frac{\big(\langle{\uparrow_1\uparrow_N}| \otimes \langle\psi^{n'}_{\text{media}}|\big) H_{\text{couple}}\big(|{\uparrow_1\downarrow_N}\rangle \otimes |\psi^{n_0}_{\text{media}}\rangle\big)}{-2h_{N\text{op}}}\Big(1 + O(\beta^{\kappa_N}) + O(\beta^{\tilde{\kappa}})\Big) \\
&+ \frac{\big(\langle{\downarrow_1\uparrow_N}| \otimes \langle\psi^{n}_{\text{media}}|\big) H_{\text{couple}}\big(|{\downarrow_1\downarrow_N}\rangle \otimes |\psi^{n'}_{\text{media}}\rangle\big)}{-2(h_{1\text{op}} - h_{N\text{op}})} \\
&\times \frac{\big(\langle{\downarrow_1\downarrow_N}| \otimes \langle\psi^{n'}_{\text{media}}|\big) H_{\text{couple}}\big(|{\uparrow_1\downarrow_N}\rangle \otimes |\psi^{n_0}_{\text{media}}\rangle\big)}{-2h_{1\text{op}}}\Big(1 + O(\beta^{\kappa_1}) + O(\beta^{\tilde{\kappa}})\Big)\Bigg].
\end{aligned} \tag{S.16}$$

We finally arrive at

$$\begin{aligned}
|\psi^{n_0,\uparrow\downarrow}_{\text{media},\downarrow\uparrow}\rangle =&\frac{J^2}{4h_{1\text{op}}h_{N\text{op}}}\Big(\gamma s_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma^2 t_{n_0}|{\downarrow_2}\rangle|\tilde{\psi}^{\uparrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle \\
&+ u_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\uparrow}_{n_0}\rangle|{\downarrow_{N-1}}\rangle + \gamma w_{n_0}|{\uparrow_2}\rangle|\tilde{\psi}^{\downarrow\downarrow}_{n_0}\rangle|{\uparrow_{N-1}}\rangle + O(\beta^{\kappa_N}) + O(\beta^{\tilde{\kappa}})\Big),
\end{aligned} \tag{S.17}$$

where κ' is defined in (16).

S.2: Calculation in the case (c)

Here, we derive the perturbed form of $|\phi^{n_0,\uparrow}_{\mathrm{tot}}\rangle$ in Eq. (63) to obtain Eq. (66). Using Eq. (S.1), the leading term of $|\phi^{n_0,\uparrow}_{\mathrm{tot},\downarrow}\rangle$ is given by

$$|\phi^{n_0,\uparrow}_{\mathrm{tot},\downarrow}\rangle = \sum_{n=1}^{2^{N-1}} |{\downarrow_1}\rangle \otimes |\phi^{n}_{\mathrm{media}}\rangle \frac{\big(\langle{\downarrow_1}| \otimes \langle\phi^{n}_{\mathrm{media}}|\big) H'_{\mathrm{couple}} \big(|{\uparrow_1}\rangle \otimes |\phi^{n_0}_{\mathrm{media}}\rangle\big)}{E'^{n_0}_{\mathrm{media}} - h_{1\mathrm{op}} - (E'^{n}_{\mathrm{media}} + h_{1\mathrm{op}})}, \tag{S.1}$$

where $h_{1\mathrm{op}} = O(\beta^{-\kappa_1})$ with $\kappa_1 \geq 1$, but $\{E'^{n}_{\mathrm{media}}\}$ are of order β^0. Then we obtain

$$|\phi^{n_0,\uparrow}_{\mathrm{tot},\downarrow}\rangle = \sum_{n=1}^{2^{N-1}} |{\downarrow_1}\rangle \otimes |\phi^{n}_{\mathrm{media}}\rangle \frac{\big(\langle{\downarrow_1}| \otimes \langle\phi^{n}_{\mathrm{media}}|\big) H'_{\mathrm{couple}} \big(|{\uparrow_1}\rangle \otimes |\phi^{n_0}_{\mathrm{media}}\rangle\big)}{-2h_{1\mathrm{op}}} \Big(1 + O(\beta^{\kappa_1})\Big). \tag{S.2}$$

We thereby obtain $|\phi^{n_0,\uparrow}_{\mathrm{tot}}\rangle$ as

$$\begin{aligned} |\phi^{n_0,\uparrow}_{\mathrm{tot}}\rangle = {} & |{\uparrow_1}\rangle \otimes |\phi^{n_0}_{\mathrm{media}}\rangle + \frac{J}{-2h_{1\mathrm{op}}} |{\downarrow_1}\rangle \otimes \Big(\gamma s'_{n_0} |{\downarrow_2}\rangle |\tilde{\phi}^{\uparrow\uparrow}_{n_0}\rangle |{\uparrow_N}\rangle + \gamma t'_{n_0} |{\downarrow_2}\rangle |\tilde{\phi}^{\uparrow\downarrow}_{n_0}\rangle |{\downarrow_N}\rangle \\ & + u'_{n_0} |{\uparrow_2}\rangle |\tilde{\phi}^{\downarrow\uparrow}_{n_0}\rangle |{\uparrow_N}\rangle + w'_{n_0} |{\uparrow_2}\rangle |\tilde{\phi}^{\downarrow\downarrow}_{n_0}\rangle |{\downarrow_N}\rangle + O(\beta^{\kappa_1})\Big). \end{aligned} \tag{S.3}$$

S.3: Proof for the existence of the maximized entanglement in three-spin systems

In this section, we prove the statement in Sec. IV. A that the maximized entanglement always exists in the systems with the Hamiltonian (81) in the high-temperature limit $\beta \to 0$, except the case of the Ising chain with $h^z_{\mathrm{media}} = 0$. We prove it by showing that the entanglement exists by letting h^z_1 and h^z_3

$$h^z_1 = h^z_3 = h_0(\beta), \tag{S.1}$$

where we assume

$$h_0(\beta) = O(\beta^{-\kappa_0}) \text{ with } \kappa_0 < -1. \tag{S.2}$$

We regard the interaction Hamiltonian as perturbation and calculate the leading order of the elements $\{F_1, F_2, p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$ of the density matrix. Then we show the inequality (8), which is a necessary and sufficient condition for the existence of the entanglement.

First, we separate the Hamiltonian (81) as follows:

$$H_{\mathrm{tot}} = \tilde{H}_{\mathrm{LO}} + \tilde{H}_{\mathrm{int}}, \tag{S.3}$$

where

$$\begin{aligned} \tilde{H}_{\mathrm{LO}} &= h^z_1 \sigma^z_1 + h^z_3 \sigma^z_3 + h^z_{\mathrm{media}} \sigma^z_2 + J^z \sigma^z_1 \sigma^z_2 + J^z \sigma^z_2 \sigma^z_3, \\ \tilde{H}_{\mathrm{int}} &= (J^x \sigma^x_1 \sigma^x_2 + J^y \sigma^y_1 \sigma^y_2) + (J^x \sigma^x_2 \sigma^x_3 + J^y \sigma^y_2 \sigma^y_3) \\ &= \Big\{ J\big[\sigma^+_1 \sigma^-_2 + \sigma^-_1 \sigma^+_2 + \gamma(\sigma^+_1 \sigma^+_2 + \sigma^-_1 \sigma^-_2)\big] + J\big[\sigma^+_2 \sigma^-_3 + \sigma^-_2 \sigma^+_3 + \gamma(\sigma^+_2 \sigma^+_3 + \sigma^-_2 \sigma^-_3)\big] \Big\}, \end{aligned} \tag{S.4}$$

h^z_{media} has an arbitrary value, $J^x = J(1+\gamma)/2$, $J^y = J(1-\gamma)/2$, and we consider $\tilde{H}_{\mathrm{int}}$ as perturbation. Because $\beta h_0(\beta) \to \infty$ as $\beta \to 0$, we only have to consider the ground state and the first excited state as the unperturbed states, which are given by

$$|{\uparrow_1 \uparrow_2 \uparrow_3}\rangle \text{ and } |{\uparrow_1 \downarrow_2 \uparrow_3}\rangle \tag{S.5}$$

with the corresponding eigenvalues

$$\epsilon_1 = -2h_0 - h^z_{\mathrm{media}} + 2J^z \text{ and } \epsilon_2 = -2h_0 + h^z_{\mathrm{media}} - 2J^z, \tag{S.6}$$

where we assume $h^z_{\mathrm{media}} > 0$, but the following discussion is also applicable to the case of $h^z_{\mathrm{media}} < 0$. Other excited states have the eigenvalues $\epsilon_1 + O(\beta^{-\kappa_0})$ and hence their thermal mixing can be ignored in the limit $\beta \to 0$.

We consider the states (S.5) up to the second-order perturbation of $\tilde{H}_{\text{int}}$:

$$k_1\biggl(|{\uparrow_1}{\uparrow_2}{\uparrow_3}\rangle - \frac{J\gamma}{2h_0 + 2h^z_{\text{media}} - 2J^z}|{\downarrow_1}{\downarrow_2}{\uparrow_3}\rangle - \frac{J\gamma}{2h_0 + 2h^z_{\text{media}} - 2J^z}|{\uparrow_1}{\downarrow_2}{\downarrow_3}\rangle + \frac{J^2\gamma}{(2h_0 - 2J^z)(2h_0 + 2h^z_{\text{media}} - 2J^z)}|{\downarrow_1}{\uparrow_2}{\downarrow_3}\rangle\biggr) \tag{S.7}$$

and

$$k_2\biggl(|{\uparrow_1}{\downarrow_2}{\uparrow_3}\rangle - \frac{J}{2h_0 - 2h^z_{\text{media}} + 2J^z}|{\downarrow_1}{\uparrow_2}{\uparrow_3}\rangle - \frac{J}{2h_0 - 2h^z_{\text{media}} + 2J^z}|{\uparrow_1}{\uparrow_2}{\downarrow_3}\rangle + \frac{J^2\gamma}{(2h_0 + 2J^z)(2h_0 - 2h^z_{\text{media}} + 2J^z)}|{\downarrow_1}{\downarrow_2}{\downarrow_3}\rangle\biggr), \tag{S.8}$$

where k_1 and k_2 are the respective normalization factors. According to the above expressions, k_1 and k_2 are of order of $1 + O(\beta^{2\kappa_0})$. By mixing these two states with the Boltzmann weights $e^{-\beta\epsilon_1}$ and $e^{-\beta\epsilon_2}$, we obtain the matrix elements $\{F_1, F_2, p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$.

Let us first consider the case $\gamma \neq 1$. In this case, the leading terms of F_1 and $\sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}}$ are given as follows, respectively:

$$\begin{aligned} F_1 &= \frac{J^2(1+\gamma^2)}{8h_0^2} + O(\beta^{1+2\kappa_0}), \\ \sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}} &= \frac{J^2\gamma}{4h_0^2} + O(\beta^{1+2\kappa_0}), \end{aligned} \tag{S.9}$$

which yields

$$F_1 - \sqrt{p_{\uparrow\uparrow}p_{\downarrow\downarrow}} = \frac{J^2(1-\gamma)^2}{8h_0^2} + O(\beta^{1+2\kappa_0}) > 0 \tag{S.10}$$

in the limit $\beta \to 0$. This gives a non-zero value of the concurrence (7) of order $\beta^{2\kappa_0}$ in the case $\gamma \neq 0$.

Next, we consider the case $\gamma = 1$. Because we assumed $J^x \geq J^y \geq J^z$, the equality $J^y = J^z = 0$ is satisfied in the case $\gamma = 1$. In this case, we have to take higher-order approximation because the first term of Eq. (S.10) vanishes. The expansions of F_2 and $\sqrt{p_{\uparrow\downarrow}p_{\downarrow\uparrow}}$ are given, respectively, by

$$\begin{aligned} F_2 &= \frac{J^2}{4h_0^2} + \frac{(h^z_{\text{media}})^2 J^2}{8}\frac{\beta}{h_0^2} - \frac{(h^z_{\text{media}})^2 J^2}{4}\frac{\beta}{h_0^3} + O(\beta^{2+3\kappa_0}), \\ \sqrt{p_{\downarrow\uparrow}p_{\uparrow\downarrow}} &= \frac{J^2}{4h_0^2} + \frac{(h^z_{\text{media}})^2 J^2}{8}\frac{\beta}{h_0^2} - \frac{(h^z_{\text{media}})^2 J^2}{2}\frac{\beta}{h_0^3} + O(\beta^{2+3\kappa_0}), \end{aligned} \tag{S.11}$$

which is followed by

$$F_2 - \sqrt{p_{\uparrow\downarrow}p_{\downarrow\uparrow}} = \frac{\beta}{4h_0^3}J^2(h^z_{\text{media}})^2 + O(\beta^{2+3\kappa_0}). \tag{S.12}$$

If the media field h^z_{media} is equal to zero, the entanglement vanishes for any values of the local fields h^z_1 and h^z_3; this is consistent with Theorem 1. For $h^z_{\text{media}} \neq 0$, the concurrence (7) is of order $\beta^{1+3\kappa_0}$ in the case $\gamma = 1$ and increases as the media field h^z_{media} is increased. We have thereby proved that the entanglement in a three-spin system with Hamiltonian (81) has always a non-zero value except the case of the Ising chain with $h^z_{\text{media}} = 0$ if we choose the local fields properly.

S.4: Calculation of the critical temperature

Here, we calculate the critical temperature analytically for the use in Sec. V. C. In order to perform the calculation, we utilize the numerically confirmed fact that the maximizing local fields take the form $h_{1\text{op}} = h_{4\text{op}} \to \infty$ in some region of the 'Symmetric' phases in Fig. 3. Let us now derive the critical temperature in the limit $h_{1\text{op}} = h_{4\text{op}} \to \infty$. In order to calculate the entanglement in this limit, we regard the coupling Hamiltonian H_{couple} between the focused spins and the media spins as perturbation. Because of the condition $h_{1\mathrm{op}} = h_{4\mathrm{op}} \to \infty$, we consider only the following four unperturbed states:

$$|{\uparrow_1\uparrow_4}\rangle \otimes |\psi_1\rangle,\ |{\uparrow_1\uparrow_4}\rangle \otimes |\psi_2\rangle,\ |{\uparrow_1\uparrow_4}\rangle \otimes |\psi_3\rangle \text{ and } |{\uparrow_1\uparrow_4}\rangle \otimes |\psi_4\rangle, \tag{S.1}$$

where $\{|\psi_n\rangle\}_{n=1}^4$ are the eigenstates of the media spins. The states $\{|\psi_n\rangle\}_{n=1}^4$ and their corresponding eigenvalues are given by the solution of the bipartite XY spin chain:

$$\begin{aligned}
|\psi_1\rangle &= \frac{1}{\sqrt{2}}\big(|{\uparrow_2\downarrow_3}\rangle - |{\downarrow_2\uparrow_3}\rangle\big),\ \epsilon_1 = -J^x - J^y,\\
|\psi_2\rangle &= \frac{1}{\sqrt{2}}\big(|{\uparrow_2\downarrow_3}\rangle + |{\downarrow_2\uparrow_3}\rangle\big),\ \epsilon_2 = J^x + J^y,\\
|\psi_3\rangle &= \mathcal{S}_-|{\downarrow_3\downarrow_4}\rangle - \mathcal{S}_+|{\uparrow_2\uparrow_3}\rangle,\ \epsilon_3 = -\sqrt{4(h^z_{\mathrm{media}})^2 + (J^x - J^y)^2},\\
|\psi_4\rangle &= \mathcal{S}_+|{\downarrow_3\downarrow_4}\rangle + \mathcal{S}_-|{\uparrow_2\uparrow_3}\rangle,\ \epsilon_4 = \sqrt{4(h^z_{\mathrm{media}})^2 + (J^x - J^y)^2},
\end{aligned} \tag{S.2}$$

where we define $\mathcal{S}_+$ and $\mathcal{S}_-$ as

$$\mathcal{S}_\pm \equiv \frac{1}{\sqrt{2}}\sqrt{1 \pm \frac{2h^z_{\mathrm{media}}}{\sqrt{4(h^z_{\mathrm{media}})^2 + (J^x - J^y)^2}}}. \tag{S.3}$$

Next, we calculate the perturbation of $\{|{\uparrow_1\uparrow_4}\rangle \otimes |\psi_n\rangle\}_{n=1}^4$ as in the proof of Theorem 2. The perturbation contribution of $|{\uparrow_1\uparrow_4}\rangle \otimes |\psi_{n_0}\rangle$ to the leading term of the element F_1 is given by

$$\frac{J^2}{4h_{1\mathrm{op}}h_{4\mathrm{op}}}\Big(\gamma s_{n_0} w_{n_0} + u_{n_0} t_{n_0} + \gamma^2 t_{n_0} u_{n_0} + \gamma w_{n_0} s_{n_0}\Big), \tag{S.4}$$

where $\{s_n, t_n, u_n, w_n\}$ are defined as the coefficients of $\{|\psi_n\rangle\}_{n=1}^4$ in (S.2):

$$|\psi_n\rangle = s_n|{\uparrow_2\uparrow_3}\rangle + t_n|{\uparrow_2\downarrow_3}\rangle + u_n|{\downarrow_2\uparrow_3}\rangle + w_n|{\downarrow_2\downarrow_3}\rangle. \tag{S.5}$$

Equations (S.4) and (S.5) correspond to Eqs. (40) and (21), respectively. Similarly, the perturbation contribution of $|{\uparrow_1\uparrow_4}\rangle \otimes |\psi_{n_0}\rangle$ to the leading term of the element F_2 is given by

$$\frac{J^2}{4h_{1\mathrm{op}}h_{4\mathrm{op}}}\big(\gamma^2 w_{n_0} s_{n_0} + \gamma u_{n_0} t_{n_0} + \gamma t_{n_0} u_{n_0} + s_{n_0} w_{n_0}\big), \tag{S.6}$$

which corresponds to Eq. (41). On the other hand, Eqs. (37)–(39) give the perturbation contribution of $|{\uparrow_1\uparrow_4}\rangle \otimes |\psi_{n_0}\rangle$ to the leading terms of the elements $\{p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$ as

$$\begin{aligned}
&\frac{J^2}{4h_{4\mathrm{op}}^2}\big(\gamma^2 s_{n_0}^2 + t_{n_0}^2 + \gamma^2 u_{n_0}^2 + w_{n_0}^2\big),\\
&\frac{J^2}{4h_{1\mathrm{op}}^2}\big(\gamma^2 s_{n_0}^2 + \gamma^2 t_{n_0}^2 + u_{n_0}^2 + w_{n_0}^2\big),\\
&\frac{J^4}{16h_{1\mathrm{op}}^2 h_{4\mathrm{op}}^2}\big(\gamma^4 s_{n_0}^2 + \gamma^2 t_{n_0}^2 + \gamma^2 u_{n_0}^2 + w_{n_0}^2\big),
\end{aligned} \tag{S.7}$$

respectively.

We have now all the ingredients for calculating the matrix elements $\{F_1, F_2, p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$ in the limit $h_{1\mathrm{op}} = h_{4\mathrm{op}} \to \infty$. We consider the mixture of the perturbed states of $|{\uparrow_1\uparrow_4}\rangle\otimes|\psi_1\rangle$, $|{\uparrow_1\uparrow_4}\rangle\otimes|\psi_2\rangle$, $|{\uparrow_1\uparrow_4}\rangle\otimes|\psi_3\rangle$ and $|{\uparrow_1\uparrow_4}\rangle\otimes|\psi_4\rangle$. These four states mix with the Boltzmann weights of $e^{-\beta(\epsilon_1+\delta\epsilon_1)}$, $e^{-\beta(\epsilon_2+\delta\epsilon_2)}$, $e^{-\beta(\epsilon_3+\delta\epsilon_3)}$ and $e^{-\beta(\epsilon_4+\delta\epsilon_4)}$, where we define the energy perturbations as $\{\delta\epsilon_i\}_{i=1}^4$. In the limit $h_{1\mathrm{op}} = h_{4\mathrm{op}} \to \infty$, we have to consider only the leading terms of $\{F_1, F_2, p_{\uparrow\uparrow}, p_{\uparrow\downarrow}, p_{\downarrow\uparrow}, p_{\downarrow\downarrow}\}$. Therefore, we ignore the energy perturbation $\{\delta\epsilon_i\}_{i=1}^4$. We thus arrive at the elements of the density matrix as

$$\begin{aligned}
Z_{\mathrm{tot}}F_1 &= \frac{J^2}{4h_0^2}\big[-4\mathcal{S}_-\mathcal{S}_+\gamma\sinh\beta|\epsilon_3| - (1+\gamma^2)\sinh\beta|\epsilon_1|\big],\\
Z_{\mathrm{tot}}F_2 &= \frac{J^2}{4h_0^2}\big[-2\mathcal{S}_-\mathcal{S}_+(1+\gamma^2)\sinh\beta|\epsilon_3| - 2\gamma\sinh\beta|\epsilon_1|\big],
\end{aligned} \tag{S.8}$$

and

$$
\begin{aligned}
Z_{\rm tot}p_{\uparrow\uparrow} &= Z_{\rm tot} = 2\cosh\beta|\epsilon_1| + 2\cosh\beta|\epsilon_3|,\\
Z_{\rm tot}p_{\uparrow\downarrow} &= \frac{J^2}{4h_0^2}\Big[(\gamma^2\mathcal{S}_-^2+\mathcal{S}_+^2)e^{-\beta\epsilon_4} + (\mathcal{S}_-^2+\gamma^2\mathcal{S}_+^2)e^{-\beta\epsilon_3} + (\gamma^2+1)\cosh\beta|\epsilon_1|\Big],\\
Z_{\rm tot}p_{\downarrow\uparrow} &= \frac{J^2}{4h_0^2}\Big[(\gamma^2\mathcal{S}_-^2+\mathcal{S}_+^2)e^{-\beta\epsilon_4} + (\mathcal{S}_-^2+\gamma^2\mathcal{S}_+^2)e^{-\beta\epsilon_3} + (\gamma^2+1)\cosh\beta|\epsilon_1|\Big],\\
Z_{\rm tot}p_{\downarrow\downarrow} &= \frac{J^4}{16h_0^4}\Big[(\gamma^4\mathcal{S}_-^2+\mathcal{S}_+^2)e^{-\beta\epsilon_4} + ((\mathcal{S}_-^2+\gamma^4\mathcal{S}_+^2)e^{-\beta\epsilon_3} + 2\gamma^2\cosh\beta|\epsilon_1|\Big],
\end{aligned}
\tag{S.9}
$$

where we defined $h_0 \equiv h_{1\rm op} = h_{4\rm op}$ with $h_0 \to \infty$.

By utilizing these parameters, we obtain a necessary and sufficient condition (8) for the existence of the entanglement in the form of the following two inequalities:

$$
\begin{aligned}
&Z_{\rm tot}^2(F_1^2 - p_{\uparrow\uparrow}p_{\downarrow\downarrow}) = \big[4\mathcal{S}_-\mathcal{S}_+\gamma\sinh\beta|\epsilon_3| + (1+\gamma^2)\sinh\beta|\epsilon_1|\big]^2\\
&- Z_{\rm tot}\big[(\gamma^4\mathcal{S}_-^2+\mathcal{S}_+^2)e^{-\beta\epsilon_4} + (\mathcal{S}_-^2+\gamma^4\mathcal{S}_+^2)e^{-\beta\epsilon_3} + 2\gamma^2\cosh\beta|\epsilon_1|\big] > 0,
\end{aligned}
\tag{S.10}
$$

or

$$
\begin{aligned}
&Z_{\rm tot}^2(F_2^2 - p_{\uparrow\downarrow}p_{\downarrow\uparrow}) = \big[2\mathcal{S}_-\mathcal{S}_+(1+\gamma^2)\sinh\beta|\epsilon_3| + 2\gamma\sinh\beta|\epsilon_1|\big]\\
&- \big[(\gamma^2\mathcal{S}_-^2+\mathcal{S}_+^2)e^{-\beta\epsilon_4} + (\mathcal{S}_-^2+\gamma^2\mathcal{S}_+^2)e^{-\beta\epsilon_3} + (\gamma^2+1)\cosh\beta|\epsilon_1|\big] > 0.
\end{aligned}
\tag{S.11}
$$